



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>6</sup> :</b> <b>G01N 33/574, 33/48, C12Q 1/68</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/54738</b> <b>(43) International Publication Date:</b> 28 October 1999 (28.10.99)
<b>(21) International Application Number:</b> PCT/US99/05766 <b>(22) International Filing Date:</b> 16 March 1999 (16.03.99)  <b>(30) Priority Data:</b> 09/061,709                      17 April 1998 (17.04.98)                      US  <b>(71) Applicant:</b> LUDWIG INSTITUTE FOR CANCER RE- SEARCH [CH/US]; 605 Third Avenue, New York, NY 10103 (US).  <b>(72) Inventors:</b> CHEN, Yao-Tseng; 525 East 68 Street, New York, NY 10021 (US). GURE, Ali; 1275 York Avenue, New York, NY 10021 (US). TSANG, Solam; 1275 York Avenue, New York, NY 10021 (US). STOCKERT, Elisabeth; 1275 York Avenue, New York, NY 10002 (US). JAGER, Elke; Steinbacher Hohl 2-28, D-60488 Frankfurt am Main (DE). KNUTH, Alexander; Steinbacher Hohl 2-28, D-60488 Frankfurt am Main (DE). OLD, Lloyd, J.; 605 Third Avenue, New York, NY 10021 (US).  <b>(74) Agent:</b> HANSON, Norman, D.; Fulbright & Jaworski LLP, 666 Fifth Avenue, New York, NY 10103 (US).		<b>(81) Designated States:</b> AU, CA, CN, JP, KR, NZ, ZA, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ISOLATED NUCLEIC ACID MOLECULE ENCODING CANCER ASSOCIATED ANTIGENS, THE ANTIGENS PER SE, AND USES THEREOF  <b>(57) Abstract</b>  The invention relates to newly identified cancer associated antigens, referred to as CT7, KOC-2 and KOC-3. The invention also relates to observations regarding known molecule KOC-1. It has been discovered that each of these molecules provokes antibodies when expressed by a subject. The ramifications of this observation are also a part of this invention.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ISOLATED NUCLEIC ACID MOLECULE ENCODING  
CANCER ASSOCIATED ANTIGENS, THE ANTIGENS PER SE,  
AND USES THEREOF

### RELATED APPLICATION

This application is a continuation in part of Serial No. 09/061,709 filed April 17, 1998, incorporated by reference.

### FIELD OF THE INVENTION

This invention relates to antigens associated with cancer, the nucleic acid molecules encoding them, as well as the uses of these.

### BACKGROUND AND PRIOR ART

It is fairly well established that many pathological conditions, such as infections, cancer, autoimmune disorders, etc., are characterized by the inappropriate expression of certain molecules. These molecules thus serve as "markers" for a particular pathological or abnormal condition. Apart from their use as diagnostic "targets", i.e., materials to be identified to diagnose these abnormal conditions, the molecules serve as reagents which can be used to generate diagnostic and/or therapeutic agents. A by no means limiting example of this is the use of cancer markers to produce antibodies specific to a particular marker. Yet another non-limiting example is the use of a peptide which complexes with an MHC molecule, to generate cytolytic T cells against abnormal cells.

Preparation of such materials, of course, presupposes a source of the reagents used to generate these. Purification from cells is one laborious, far from sure method of doing so. Another preferred method is the isolation of nucleic acid molecules which encode a particular marker, followed by the use of the isolated encoding molecule to express the desired molecule.

Two basic strategies have been employed for the detection of such antigens, in e.g., human tumors. These will be referred to as the genetic approach and the biochemical approach. The genetic approach is exemplified by, e.g., dePlaen et al., Proc. Natl. Sci. USA 85: 2275

(1988), incorporated by reference. In this approach, several hundred pools of plasmids of a cDNA library obtained from a tumor are transfected into recipient cells, such as COS cells, or into antigen-negative variants of tumor cell lines which are tested for the expression of the specific antigen. The biochemical approach, exemplified by, e.g., O. Mandelboim, et al., *Nature* 369: 69 (1994) incorporated by reference, is based on acidic elution of peptides which have bound to MHC-class I molecules of tumor cells, followed by reversed-phase high performance liquid chromatography (HPLC). Antigenic peptides are identified after they bind to empty MHC-class I molecules of mutant cell lines, defective in antigen processing, and induce specific reactions with cytotoxic T-lymphocytes. These reactions include induction of CTL proliferation, TNF release, and lysis of target cells, measurable in an MTT assay, or a  $^{51}\text{Cr}$  release assay.

These two approaches to the molecular definition of antigens have the following disadvantages: first, they are enormously cumbersome, time-consuming and expensive; and second, they depend on the establishment of cytotoxic T cell lines (CTLs) with predefined specificity.

The problems inherent to the two known approaches for the identification and molecular definition of antigens is best demonstrated by the fact that both methods have, so far, succeeded in defining only very few new antigens in human tumors. See, e.g., van der Bruggen et al., *Science* 254: 1643-1647 (1991); Brichard et al., *J. Exp. Med.* 178: 489-495 (1993); Coulie, et al., *J. Exp. Med.* 180: 35-42 (1994); Kawakami, et al., *Proc. Natl. Acad. Sci. USA* 91: 3515-3519 (1994).

Further, the methodologies described rely on the availability of established, permanent cell lines of the cancer type under consideration. It is very difficult to establish cell lines from certain cancer types, as is shown by, e.g., Oettgen, et al., *Immunol. Allerg. Clin. North. Am.*

10: 607-637 (1990). It is also known that some epithelial cell type cancers are poorly susceptible to CTLs in vitro, precluding routine analysis. These problems have stimulated the art to develop additional methodologies for identifying cancer associated antigens.

One key methodology is described by Sahin, et al., Proc. Natl. Acad. Sci. USA 92: 11810-11913 (1995), incorporated by reference. Also, see U.S. Patent No. 5,698,396, and Application Serial No. 08/479,328, filed on June 7, 1995 and January 3, 1996, respectively. All three of these references are incorporated by reference. To summarize, the method involves the expression of cDNA libraries in a prokaryotic host. (The libraries are secured from a tumor sample). The expressed libraries are then immunoscreened with absorbed and diluted sera, in order to detect those antigens which elicit high titer humoral responses. This methodology is known as the SEREX method ("Serological identification of antigens by Recombinant Expression Cloning"). The methodology has been employed to confirm expression of previously identified tumor associated antigens, as well as to detect new ones. See the above referenced patent applications and Sahin, et al., supra, as well as Crew, et al., EMBO J 144: 2333-2340 (1995).

This methodology has been applied to a range of tumor types, including those described by Sahin et al., supra, and Pfreundschuh, supra, as well as to esophageal cancer (Chen et al., Proc. Natl. Acad. Sci. USA 94: 1914-1918 (1997)); lung cancer (Güre et al., Cancer Res. 58: 1034-1041 (1998)); colon cancer (Serial No. 08/948, 705 filed October 10, 1997) incorporated by reference, and so forth. Among the antigens identified via SEREX are the SSX2 molecule (Sahin et al., Proc. Natl. Acad. Sci. USA 92: 11810-11813 (1995); Tureci et al., Cancer Res. 56: 4766-4772 (1996); NY-ESO-1 Chen, et al., Proc. Natl. Acad. Sci. USA 94: 1914-1918 (1997); and SCP1 (Serial No. 08/892,705 filed July 15, 1997) incorporated by reference.

Analysis of SEREX identified antigens has shown overlap between SEREX defined and CTL defined antigens. MAGE-1, tyrosinase, and NY-ESO-1 have all been shown to be recognized by patient antibodies as well as CTLs, showing that humoral and cell mediated responses do act in concert.

It is clear from this summary that identification of relevant antigens via SEREX is a desirable aim. The inventors have modified standard SEREX protocols and have screened a cell line known to be a good source of the antigens listed supra, using allogeneic patient sample. New antigens have been identified in this way and have been studied. Also, a previously known molecule has now been identified via SEREX techniques.

#### DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

##### Example 1

The melanoma cell referred to as SK-MEL-37 was used, because it has been shown to express a number of members of the CT antigen family, including MAGE-1 (Chen et al., Proc. Natl. Acad. Sci. USA 91: 1004-1008 (1994); NY-ESO-1 (Chen et al. Proc. Natl. Acad. Sci. USA 94: 1914-1918 (1997)); and various members of the SSX family (Gure et al., Int. J. Cancer 72: 965-971 (1997)).

Total RNA was extracted from cultured samples of SK-MEL-37 using standard methods, and this was then used to construct a cDNA library in commercially available,  $\lambda$ ZAP expression vector, following protocols provided by the manufacturer. The cDNA was then transfected into E. coli and screened, following Sahin et al., Proc. Natl. Acad. Sci. USA 92: 11810-11813 (1995), incorporated by reference, and Pfreundschuh, U.S. Patent No. 5,698,396, also incorporated by reference. The screening was done with allogeneic patient serum "NW38." This serum had been shown, previously, to contain high titer antibodies against MAGE-1 and NY-

ESO-1. See, e.g., Jäger et al., J. Exp. Med. 187: 265-270 (1998), incorporated by reference. In brief, serum was diluted 1:10, preabsorbed with lysates of transfected E. coli, further diluted to 1:2000, and then incubated overnight at room temperature with nitrocellulose membranes containing phage plaques, prepared in accordance with Sahin et al., and Pfreundschuh, supra. The library contained a total of  $2.3 \times 10^7$  primary clones. After washing, the filters were incubated with alkaline phosphatase conjugated, goat anti-human Fc $\gamma$  secondary antibodies, and were then visualized by incubating with 5-bromo-4-chloro-3-indolyl phosphate, and nitroblue tetrazolium.

After screening  $1.5 \times 10^5$  of the clones, a total of sixty-one positives had been identified. Given this number, screening was stopped, and the positive clones were subjected to further analysis.

### Example 2

The positive clones identified in example 1, supra, were purified, the inserts were excised in vitro, and inserted into a commercially available plasmid, pBK-CMV, and then evaluated on the basis of restriction mapping with EcoRI and XbaI. Clones which represented different inserts on the basis of this step were sequenced, using standard methodologies.

There was a group of 10 clones, which could not be classified other than as “miscellaneous genes”, in that they did not seem to belong to any particular family. They consisted of 9 distinct genes, of which four were known, and five were new. The fifty one remaining clones were classified into four groups. The data are presented in Tables 1 and 2, which follow.

The largest group are genes related to KOC (“KH-domain containing gene, overexpressed in cancer” which has been shown to be overexpressed in pancreatic cancer, and maps to



chromosome 7p11.5. See Müller-Pillasch et al., *Oncogene* 14: 2729-2733 (1997). Two of the 33 were derived from the KOC gene, and the other 31 were derived from two previously unidentified, but related genes. Examples 6 et seq. describe work on this group of clones.

Eleven clones, i.e., Group 2, were MAGE sequences. Four were derived from MAGE-4a, taught by DePlaen et al., *Immunogenetics* 40: 360-369, Genbank U10687, while the other 7 hybridized to a MAGE-4a probe, derived from the 5' sequence, suggesting they belong to the MAGE family.

The third group consisted of five clones of the NY-ESO-1 family. Two were identical to the gene described by Chen et al., *Proc. Natl. Acad. Sci. USA* 94: 1914-1918 (1997), and in Serial No. 08/725,182, filed October 3, 1996, incorporated by reference. The other three were derived from a second member of the NY-ESO-1 family, i.e., LAGE-1. See U.S. application Serial No. 08/791,495, filed January 27, 1997 and incorporated by reference.

The fourth, and final group, related to a novel gene referred to as CT7. This gene, the sequence of which is presented as SEQ ID NO: 1, was studied further.

Table 1. SEREX-identified genes from allogeneic screening of SK-MEL-37 library

Gene group	# of clones	Comments
KOC	33	derived from 3 related genes
MAGE	11	predominantly MAGE-4a (see text)
NY-ESO-1	5	derived from 2 related genes (NY-ESO-1, LAGE-1)
CT7	2	new cancer/testis antigen
Miscellaneous	10	see Table 2

Table 2. SEREX-identified genes from allogeneic screening of SK-MEL-37 library--  
Miscellaneous group

Clone designation	Gene
MNW-4, MNW-7	S-adenyl homocysteine hydrolase
MNE-6a	Glutathione synthetase
MNW-24	proliferation-associated protein p38-2G4
MNW-27a	phosphoribosyl pyrophosphate synthetase-associated protein 39
MNW-6b	unknown gene, identical to sequence tags from pancreas, uterus etc.
MNW-14b	unknown gene, identical to sequence tags from lung, brain, fibroblast etc.
MNW-34a	unknown gene, identical to sequence tags from multiple tissues
MNW-17	unknown gene, identical to sequence tags from pancreas and fetus
MNW-29a	unknown gene, no significant sequence homology, universally expressed

### Example 3

The two clones for CT7, referred to supra, were 2184 and 1965 base pairs long. Analysis of the longer one was carried out. It presented an open reading frame of 543 amino acids, which extended to the 5' end of the sequence, indicating that it was a partial cDNA clone.

In order to identify the complete sequence, and to try to identify additional, related genes, a human testicular cDNA library was prepared, following standard methods, and screened with probes derived from the longer sequence, following standard methods.

Eleven positives were detected, and sequenced, and it was found that all derived from the same gene. When the polyA tail was excluded, full length transcript, as per SEQ ID NO: 1, consisted of 4265 nucleotides, broken down into 286 base pairs of untranslated 5' - region, a coding region of 3429 base pairs, and 550 base pairs of untranslated 3' region. The predicted protein is 1142 amino acids long, and has a calculated molecular mass of about 125 kilodaltons. See SEQ ID NO: 2.

The nucleotide and deduced amino acid sequences were screened against known databases, and there was some homology with the MAGE-10 gene, described by DePlaen et al., Immunogenetics 40: 360-369 (1994). The homology was limited to about 210 carboxy terminal amino acids, i.e., amino acids 908-1115 of the subject sequence, and 134-342 of MAGE-10. The percent homology was 56%, rising to 75% when conservative changes are included.

There was also extensive homology with a sequence reported by Lucas et al., Canc. Res. 58: 743-752 (1998), and application Serial No. 08/845,528 filed April 25, 1997, also incorporated by reference. A total of 14 nucleotides differ in the open reading frame, resulting in a total of 11 amino acids which differ between the sequences.

The 5' region of the nucleotide and sequence and corresponding amino acid sequence demonstrates a strikingly repetitive pattern, with repeats rich in serine, proline, glutamine, and leucine, with an almost invariable core of PQSPLQI (SEQ ID NO: 3). In the middle of the molecule, 11 almost exact repeats of 35 amino acids were observed. The repetitive portions make up about 70% of the entire sequence, begin shortly after translation initiation, at position 15, and ending shortly before the region homologous to MAGE 4a.

#### Example 4

The expression pattern for mRNA of CT7 was then studied, in both normal and malignant tissues. RT-PCR was used, employing primers specific for the gene. The estimated melting temperature of the primers was 65-70°C, and they were designed to amplify 300-600 base pair segments. A total of 35 amplification cycles were carried out, at an annealing temperature of 60°C. Table 3, which follows, presents the data for human tumor tissues. CT7 was expressed in a number of different samples. Of fourteen normal tissues tested, there was strong expression in testis, and none in colon, brain, adrenal, lung, breast, pancreas, prostate, thymus or uterus tissue. There was low level expression in liver, kidney, placenta and fetal brain, with fetal brain showing three transcripts of different size. The level of expression was at least 20-50 times lower than in testis. Melanoma cell lines were also screened. Of these 7 of the 12 tested showed strong expression, and one showed weak expression.

Table 3. CT7 mRNA expression in various humor tumors by RT-PCR

Tumor type	mRNA, positive/total
Melanoma	7/10
Breast cancer	3/10
Lung cancer	3/9
Head/neck cancer	5/14
Bladder cancer	4/9
Colon cancer	1/10
Leimyosarcoma	1/4
synovial sarcoma	2/4
Total	26/70

#### Example 5

Southern blotting experiments were then carried out to determine if CT7 belonged to a family of genes. In these experiments, genomic DNA was extracted from normal human tissues. It was digested with BamHI, EcoRI, and HindIII, separated on a 0.7% agarose gel, blotted onto a nitrocellulose filter, and hybridized, at high stringency (65°C, aqueous buffer), with a <sup>32</sup>P labelled probe, derived from SEQ ID NO: 1.

The blotting showed anywhere from two to four bands, suggesting one or two genes in the family.

#### Example 6

As noted in example 2, supra, thirty three of the sixty one positive clones were related to KOC. Clones were sequenced using standard methodologies. As indicated supra, one clone

was identical to KOC, initially reported by Müller-Pillasch, et al., supra. Given that two additional related sequences were identified, the known KOC gene is referred to as KOC-1 hereafter (SEQ ID NO: 4). The second clone, referred to as KOC-2 hereafter, was found once. The sequence is presented as SEQ ID NO: 5. Its deduced amino acid sequence is 72.5% identical to that for KOC-1.

The third sequence, KOC-3, appeared thirty times (SEQ ID NO: 6). Its deduced amino acid sequence is 63% identical to KOC-1.

Testicular cDNA libraries were analyzed in the same way that the SK-MEL-37 library was analyzed, i.e., with allogeneic serum from NW-38. See example 3, supra.

Following analysis of testicular libraries, a longer form of KOC-2 was isolated. This is presented as SEQ ID NO: 7. When SEQ ID NOS: 5 & 7 are compared, the former is 1705 base pairs in length, without a polyA tail. It contains 1362 base pairs of coding sequence, and 343 base pairs of 3' untranslated sequence. Nucleotides 275-1942 of SEQ ID NO: 7 are identical to nucleotides 38-1705 of SEQ ID NO: 5.

The sequence of KOC-3, set forth as SEQ ID NO: 6, is 3412 base pairs long, and consists of 72 base pairs of 5' untranslated region, 1707 base pairs of open reading frame, and 1543 base pairs of untranslated, 3' region. An alternate form was also isolated, (SEQ ID NO: 8), and is 129 base pairs shorter than SEQ ID NO: 6.

#### Example 7

Expression patterns for KOC-1, KOC-2 and KOC-3 were then studied, using RT-PCR and the following primer pairs:

GAAAGTATCT TCAAGGACGC C

CTGCAAGGGG TTTTGCTGGG CG

(SEQ ID NOS: 9 & 10).

TCCTTGCGCG CTGCGGCCTC AG

CCAACTGGTG GCCATTCAGCT TC

(SEQ ID NOS: 11 & 12)

GCTCTTTGGG GACAGGAAGG TC

GACGTTGACA ACGGCGGTTT CT

(SEQ ID NOS: 13 & 14).

SEQ ID NOS: 9 & 10 were designed to amplify KOC-1 while SEQ ID NOS: 11 & 12 were designed to amplify KOC-2, and SEQ ID NOS: 13 & 14 were designed to amplify KOC-3.

To carry out the RT-PCR, relevant primer pairs were added to cDNA samples prepared from various mRNAs by reverse transcription. PCR was then carried out at an annealing temperature of 60°C, and extension at 72°C, for 35 cycles. The resulting products were then analyzed by gel electrophoresis.

SEQ ID NOS 9 & 10 amplify nucleotides 305-748 of SEQ ID NO: 1. A variety of normal and malignant cell types were tested. Strong expression was found in testis, moderate expression in normal brain, and low levels of expression were found in normal colon, kidney, and liver.

The Müller-Pillasch paper, cited supra, identified expression of KOC-1 in pancreatic tumor cell lines, gastric cancer, and normal placenta, via Northern blotting. This paper also reported that normal heart, brain, lung, liver, kidney and pancreatic tissue were negative for

KOC-1 expression. The difference in results suggests that the level of expression of KOC-1 is very low in normal tissues.

When KOC-2 expression was studied, the only positive normal tissue was testis (brain, liver, kidney and colon were negative).

Modification of the protocol for detecting KOC-2 resulted in positives in normal kidney, liver and melanoma.

When KOC-3 expression was studied, it was found that the gene was universally expressed in normal tissues, with highest expression in testis.

The pattern of expression of KOC-3 in different melanoma cell lines was analyzed, using standard Northern blotting. Over expression in several cell lines was observed, which is consistent with the more frequent isolation of this clone than any other.

### Example 8

A study was carried out to determine if KOC-1 is expressed at higher levels in melanoma cells, as compared to normal skin cells. This was done using representational difference analysis, or "RDA." See Lisitsyn, et al. Science 259: 946-951 (1993), and O'Neill, et al. Nucl. Acids Res. 25:2681-2 (1997), both of which are incorporated by reference. Specifically, tester cDNA was taken from SK-MEL-37, and driver cDNA was taken from a skin sample representing mRNA from various cell types in the skin. The cDNAs were digested with either Tsp509I, Hsp92II, or DpnII. When DpnII was the enzyme used for digestion, adaptor oligonucleotides R-Bgl-24, J-Bgl-24, and N-Bgl-24 described by O'Neill, et al., supra, and Hubank, et al. Nucl. Acids Res.



22:5640-5648 (1994) were used. When Tsp509I was the endonuclease, the same adaptors were used, as were R-Tsp-12, i.e.:

AATTTGCGGT GA

(SEQ ID NO: 15)

J-Tsp-12, i.e.:

AATTTGTTCA TG

(SEQ ID NO: 16)

and N-Tsp-12, i.e.:

AATTTTCCCT CG

(SEQ ID NO: 17)

When Hsp92II was the endonuclease, the adaptors were:

R-Hsp-24, i.e.:

AGCACTCTCC AGCCTCTCAC CATG

(SEQ ID NO: 18);

J-Hsp-24, i.e.:

ACCGACGTCG ACTATCATG CATG

(SEQ ID NO: 19);

N-Hsp-24, i.e.:

AGGCAACTGT GCTATCCGAG CATG

(SEQ ID NO: 20);

R-Hsp-8, i.e.:

GTGAGAGG

(SEQ ID NO: 21);

J-Hsp-8, i.e.:

CATGGATG

(SEQ ID NO: 22);

N-Hsp-8, i.e.:

CTCGGATA

(SEQ ID NO: 23).

In order to hybridize tester and driver, either 3XEE buffer (30mM EPPS, pH8, 3mM EDTA), or a buffer of 2.4M tetraethylammonium chloride (TEACl) 3mM EDTA, 10mM Tris HC1, pH8, was used. When DNA was dissolved in 10  $\mu$ l of TEACl buffer, it was denatured at 80°C for 10 minutes, followed by renaturing at 42°C for 20 hours. Amplicons were gel purified, and the DP3 or DP2 product was ligated into BamHI (when DpnII was used), EcoRI (when Tsp 509I was used), or SpHI (when Hsp92II was used), cloning vectors were digested, and then sequenced. Sequence analysis of the cDNA molecules derived from these experiments identified KOC-1 as one of the genes isolated, indicating that KOC-1 mRNA is present at a higher level in Sk-Mel 37 cells as compared to normal skin cells.

The foregoing examples describe the isolation of a nucleic acid molecule which encodes a cancer associated antigen. "Associated" is used herein because while it is clear that the relevant molecule was expressed by several types of cancer, other cancers, not screened herein, may also express the antigen.

The invention relates to those nucleic acid molecules which encode the antigens CT7, KOC-2 and KOC-3 , as described herein, such as a nucleic acid molecule consisting of the nucleotide sequence SEQ ID NO: 1, molecules comprising the nucleotide sequence of SEQ ID

NO: 5, 6, 7 or 8 and so forth. Also embraced are those molecules which are not identical to SEQ ID NOS: 1, 5, 6, 7 or 8, but which encode the same antigen.

Also a part of the invention are expression vectors which incorporate the nucleic acid molecules of the invention, in operable linkage (i.e., "operably linked") to a promoter. Construction of such vectors, such as viral (e.g., adenovirus or Vaccinia virus) or attenuated viral vectors is well within the skill of the art, as is the transformation or transfection of cells, to produce eukaryotic cell lines, or prokaryotic cell strains which encode the molecule of interest. Exemplary of the host cells which can be employed in this fashion are COS cells, CHO cells, yeast cells, insect cells (e.g., Spodoptera frugiperda), NIH 3T3 cells, and so forth. Prokaryotic cells, such as E. coli and other bacteria may also be used. Any of these cells can also be transformed or transfected with further nucleic acid molecules, such as those encoding cytokines, e.g., interleukins such as IL-2, 4, 6, or 12 or HLA or MHC molecules.

Also a part of the invention are the antigens described herein, both in original form and in any different post translational modified forms. The molecules are large enough to be antigenic without any posttranslational modification, and hence are useful as immunogens, when combined with an adjuvant (or without it), in both precursor and post-translationally modified forms. Antibodies produced using these antigens, both poly and monoclonal, are also a part of the invention as well as hybridomas which make monoclonal antibodies to the antigens. The whole protein can be used therapeutically, or in portions, as discussed infra. Also a part of the invention are antibodies against this antigen, be these polyclonal, monoclonal, reactive fragments, such as Fab, (F(ab)<sub>2</sub>)' and other fragments, as well as chimeras, humanized antibodies, recombinantly produced antibodies, and so forth.

As is clear from the disclosure, one may use the proteins and nucleic acid molecules of the invention diagnostically. The SEREX methodology discussed herein is premised on an immune response to a pathology associated antigen. Hence, one may assay for the relevant pathology via, e.g., testing a body fluid sample of a subject, such as serum, for reactivity with the antigen per se. Reactivity would be deemed indicative of possible presence of the pathology. So, too, could one assay for the expression of any of the antigens via any of the standard nucleic acid hybridization assays which are well known to the art, and need not be elaborated upon herein. One could assay for antibodies against the subject molecules, using standard immunoassays as well.

Analysis of SEQ ID NO: 1, 5, 6, 7 and 8 will show that there are 5' and 3' non-coding regions presented therein. The invention relates to those isolated nucleic acid molecules which contain at least the coding segment, i.e., nucleotides 54-593, of SEQ ID NO: 1, nucleotides 1-1019 of SEQ ID NO: 3, nucleotides 73-1780 of SEQ ID NO: 8, and so forth, and which may contain any or all of the non-coding 5' and 3' portions.

Also a part of the invention are portions of the relevant nucleic acid molecules which can be used, for example, as oligonucleotide primers and/or probes, such as one or more of SEQ ID NOS: 7, 8, 9, 10, 11, 12, 13 or 14 as well as amplification product like nucleic acid molecules comprising at least nucleotides 305-748 of SEQ ID NO: 1.

As was discussed supra, study of other members of the "CT" family reveals that these are also processed to peptides which provoke lysis by cytolytic T cells. There has been a great deal of work on motifs for various MHC or HLA molecules, which is applicable here. Hence, a further aspect of the invention is a therapeutic method, wherein one or more peptides derived

from the antigens of the invention which bind to an HLA molecule on the surface of a patient's tumor cells are administered to the patient, in an amount sufficient for the peptides to bind to the MHC/HLA molecules, and provoke lysis by T cells. Any combination of peptides may be used. These peptides, which may be used alone or in combination, as well as the entire protein or immunoreactive portions thereof, may be administered to a subject in need thereof, using any of the standard types of administration, such as intravenous, intradermal, subcutaneous, oral, rectal, and transdermal administration. Standard pharmaceutical carriers, adjuvants, such as saponins, GM-CSF, and interleukins and so forth may also be used. Further, these peptides and proteins may be formulated into vaccines with the listed material, as may dendritic cells, or other cells which present relevant MHC/peptide complexes.

Similarly, the invention contemplates therapies wherein nucleic acid molecules which encode the proteins of the invention, one or more or peptides which are derived from these proteins are incorporated into a vector, such as a Vaccinia or adenovirus based vector, to render it transfectable into eukaryotic cells, such as human cells. Similarly, nucleic acid molecules which encode one or more of the peptides may be incorporated into these vectors, which are then the major constituent of nucleic acid bases therapies.

Any of these assays can also be used in progression/regression studies. One can monitor the course of abnormality involving expression of these antigens simply by monitoring levels of the protein, its expression, antibodies against it and so forth using any or all of the methods set forth supra.

It should be clear that these methodologies may also be used to track the efficacy of a therapeutic regime. Essentially, one can take a baseline value for a protein of interest using any

of the assays discussed supra, administer a given therapeutic agent, and then monitor levels of the protein thereafter, observing changes in antigen levels as indicia of the efficacy of the regime.

As was indicated supra, the invention involves, inter alia, the recognition of an "integrated" immune response to the molecules of the invention. One ramification of this is the ability to monitor the course of cancer therapy. In this method, which is a part of the invention, a subject in need of the therapy receives a vaccination of a type described herein. Such a vaccination results, e.g., in a T cell response against cells presenting HLA/peptide complexes on their cells. The response also includes an antibody response, possibly a result of the release of antibody provoking proteins via the lysis of cells by the T cells. Hence, one can monitor the effect of a vaccine, by monitoring an antibody response. As is indicated, supra, an increase in antibody titer may be taken as an indicia of progress with a vaccine, and vice versa. Hence, a further aspect of the invention is a method for monitoring efficacy of a vaccine, following administration thereof, by determining levels of antibodies in the subject which are specific for the vaccine itself, or a large molecule of which the vaccine is a part.

The identification of the subject proteins as being implicated in pathological conditions such as cancer also suggests a number of therapeutic approaches in addition to those discussed supra. The experiments set forth supra establish that antibodies are produced in response to expression of the protein. Hence, a further embodiment of the invention is the treatment of conditions which are characterized by aberrant or abnormal levels of one or more of the proteins, via administration of antibodies, such as humanized antibodies, antibody fragments, and so forth. These may be tagged or labelled with appropriate cystostatic or cytotoxic reagents.

T cells may also be administered. It is to be noted that the T cells may be elicited in vitro using immune responsive cells such as dendritic cells, lymphocytes, or any other immune responsive cells, and then reperfused into the subject being treated.

Note that the generation of T cells and/or antibodies can also be accomplished by administering cells, preferably treated to be rendered non-proliferative, which present relevant T cell or B cell epitopes for response, such as the epitopes discussed supra.

The therapeutic approaches may also include antisense therapies, wherein an antisense molecule, preferably from 10 to 100 nucleotides in length, is administered to the subject either "neat" or in a carrier, such as a liposome, to facilitate incorporation into a cell, followed by inhibition of expression of the protein. Such antisense sequences may also be incorporated into appropriate vaccines, such as in viral vectors (e.g., Vaccinia), bacterial constructs, such as variants of the known BCG vaccine, and so forth.

Also a part of the inventions are Peptides, such as those set forth in Figure 1, and those which have as a core sequence

PQSPLQI (SEQ ID NO.: 3)

These peptides may be used therapeutically, via administration to a patient who expresses CT7 in connection with a pathology, as well as diagnostically, i.e., to determine if relevant antibodies are present and so forth.

Other features and applications of the invention will be clear to the skilled artisan, and need not be set forth herein. The terms and expression which have been employed are used as terms of description and not of limitation, and there is no intention in the use of such terms and expression of excluding any equivalents of the features shown and described or portions thereof, it being recognized that various modifications are possible within the scope of the invention.

We claim:

1. Isolated nucleic acid molecule which encodes a cancer associated antigen, whose amino acid sequence is identical to the amino sequence encoded by nucleotides 287 to 3714 of SEQ ID NO: 1.
2. The isolated nucleic acid molecule of claim 1, consisting of nucleotides 287-3714 of SEQ ID NO: 1.
3. The isolated nucleic acid molecule of claim 1, consisting of anywhere from nucleotide 1 through nucleotide 4265 of SEQ ID NO: 1, with the proviso that said isolated nucleic acid molecule contains at least nucleotides 287-3714 of SEQ ID NO: 1.
4. Expression vector comprising the isolated nucleic acid molecule of claim 1, operably linked to a promoter.
5. Expression vector comprising the isolated nucleic acid molecule of claim 3, operably linked to a promoter.
6. Eukaryotic cell line or prokaryotic cell strain, transformed or transfected with the expression vector of claim 4.
7. Eukaryotic cell line or prokaryotic cell strain, transformed or transfected with the expression vector of claim 5.
8. Isolated cancer associated antigen comprising all or part of the amino acid sequence encoded by nucleotides 287-3714 of SEQ ID NO: 1.
9. Eukaryotic cell line or prokaryote cell strain, transformed or transfected with the isolated nucleic acid molecule of claim 1.
10. The eukaryotic cell line of claim 9, wherein said cell line is also transfected with a nucleic acid molecule coding for a cytokine.



11. The eukaryotic cell line of claim 10, wherein said cell line is further transfected by a nucleic acid molecule coding for an HLA molecule.
12. The eukaryotic cell line of claim 10, wherein said cytokine is an interleukin.
13. The biologically pure culture of claim 12, wherein said interleukin is IL-2, IL-4 or IL-12.
14. The eukaryotic cell line of claim 9, wherein said cell line has been rendered non-proliferative.
15. The eukaryotic cell line of claim 9, wherein said cell line is a fibroblast cell line.
16. Expression vector comprising a mutated or attenuated virus and the isolated nucleic acid molecule of claim 1.
17. The expression vector of claim 16, wherein said virus is adenovirus or vaccinia virus.
18. The expression vector of claim 17, wherein said virus is vaccinia virus.
19. The expression vector of claim 17, wherein said virus is adenovirus.
20. Expression system useful in transfecting a cell, comprising (i) a first vector containing a nucleic acid molecule which codes for the isolated cancer associated antigen of claim 8 and (ii) a second vector selected from the group consisting of (a) a vector containing a nucleic acid molecule which codes for an MHC or HLA molecule which presents an antigen derived from said cancer associated antigen and (b) a vector containing a nucleic acid molecule which codes for an interleukin.
21. Isolated cancer associated antigen comprising the amino acid sequence encoded by nucleotides 287-3714 of SEQ ID NO: 1.

22. Immunogenic composition comprising the isolated antigen of claim 21, and a pharmaceutically acceptable adjuvant.

23. The immunogenic composition of claim 22, wherein said adjuvant is a cytokine, a saponin, or GM-CSF.

24. Immunogenic composition comprising at least one peptide consisting of an amino acid sequence of from 8 to 12 amino acids concatenated to each other in the isolated cancer associated antigen of claim 21, and a pharmaceutically acceptable adjuvant.

25. The immunogenic composition of claim 24, wherein said adjuvant is a saponin, a cytokine, or GM-CSF.

26. The immunogenic composition of claim 24, wherein said composition comprises a plurality of peptides which complex with a specific MHC molecule.

27. Isolated peptide derived from the amino acid sequence encoded by SEQ ID NO: 1, wherein said isolated peptide binds to an HLA molecule, is a nonamer, decamer or undecamer, and comprises the amino acid sequence of SEQ ID NO: 3, from one to three additional N-terminal amino acid, and up to four additional C terminal amino acids.

28. Immunogenic composition which comprises at least one expression vector which encodes for a peptide derived from the amino acid sequence encoded by SEQ ID NO: 1, and an adjuvant or carrier.

29. The immunogenic composition of claim 28, wherein said at least one expression vector codes for a plurality of peptides.

30. Vaccine useful in treating a subject afflicted with a cancerous condition comprising the isolated cell line of claim 11 and a pharmacologically acceptable adjuvant.

31. The vaccine of claim 30, wherein said cell line has been rendered non-proliferative.
32. The vaccine of claim 31, wherein said cell line is a human cell line.
33. A composition of matter useful in treating a cancerous condition comprising a non proliferative cell line having expressed on its surface a peptide derived from the amino acid sequence encoded by SEQ ID NO: 1.
34. The composition of matter of claim 33, wherein said cell line is a human cell line.
35. A composition of matter useful in treating a cancerous condition, comprising (i) a peptide derived from the amino acid sequence encoded by SEQ ID NO: 1, (ii) an MHC or HLA molecule, and (iii) a pharmaceutically acceptable carrier.
36. Isolated antibody which is specific for the antigen of claim 21.
37. The isolated antibody of claim 36, wherein said antibody is a monoclonal antibody.
38. Method for screening for cancer in a sample, comprising contacting said sample with a nucleic acid molecule which hybridizes to all or part of SEQ ID NO: 1, and determining hybridization as an indication of cancer cells in said sample.
39. A method for screening for cancer in a sample, comprising contacting said sample with the isolated antibody of claim 36, and determining binding of said antibody to a target as an indicator of cancer.
40. Method for diagnosing a cancerous condition in a subject, comprising contacting an immune reactive cell containing sample of said subject to a cell line transfected with the isolated nucleic acid molecule of claim 1, and determining interaction of said transfected cell line with said immunoreactive cell, said interaction being indicative of said cancer condition.

41. A method for determining regression, progression of onset of a cancerous condition comprising monitoring a sample from a patient with said cancerous condition for a parameter selected from the group consisting of (i) CT7 protein, (ii) a peptide derived from CT7 protein (iii) cytolytic T cells specific for said peptide and an MHC molecule with which it non-covalently complexes, and (iv) antibodies specific for said CT7 protein, wherein amount of said parameter is indicative of progression or regression or onset of said cancerous condition.

42. Method of claim 41, wherein said sample is a body fluid or exudate.

43. Method of claim 41, wherein said sample is a tissue.

44. Method of claim 41, comprising contacting said sample with an antibody which specifically binds with said protein or peptide.

45. Method of claim 44, wherein said antibody is labelled with a radioactive label or an enzyme.

46. Method of claim 44, wherein said antibody is a monoclonal antibody.

47. Method of claim 41, comprising amplifying RNA which codes for said protein.

48. Method of claim 47, wherein said amplifying comprises carrying out polymerase chain reaction.

49. Method of claim 41, comprising contacting said sample with a nucleic acid molecule which specifically hybridizes to a nucleic acid molecule which codes for or expresses said protein.

50. Method of claim 41, comprising assaying said sample for shed protein.

51. Method of claim 41, comprising assaying said sample for antibodies specific for said CT7 protein, by contacting said sample with CT7 protein.

52. Method for diagnosing a cancerous condition comprising assaying a sample taken from a subject for an immunoreactive cell specific for a peptide derived from CT7, complexed to an MHC molecule, presence of said immunoreactive cell being indicative of said cancerous condition.

53. An isolated nucleic acid molecule which encodes a protein and which has a complementary sequence which hybridizes, under stringent conditions, to at least one of the nucleotide sequences set forth at SEQ ID NO: 5, 6, 7 or 8.

54. The isolated nucleic acid molecule of claim 53, wherein said protein is the protein encoded by the nucleotide sequence of SEQ ID NO: 5, 6, 7 or 8.

55. The isolated nucleic acid molecule of claim 53, selected from the group consisting of nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO: 5, 6, 7 or 8.

56. Expression vector comprising the isolated nucleic acid molecule of claim 54, operably linked to a promoter.

57. Expression vector comprising the isolated nucleic acid molecule of claim 55, operably linked to a promoter.

58. Recombinant cell comprising the expression vector of claim 56.

59. Recombinant cell comprising the expression vector of claim 57.

60. Recombinant cell comprising the isolated nucleic acid molecule of claim 54.

61. Recombinant cell comprising the isolated nucleic acid molecule of claim 55.

62. Recombinant cell of claim 58, further comprising an expression vector which contains a nucleic acid molecule encoding a cytokine, operably linked to a promoter.

63. Recombinant cell of claim 59, further comprising an expression vector which contains a nucleic acid molecule encoding a cytokine, operably linked to a promoter.

64. Recombinant cell of claim 60, further comprising a nucleic acid molecule which encodes a cytokine.

65. Recombinant cell of claim 61, further comprising a nucleic acid molecule which encodes a cytokine.

66. The recombinant cell of claim 62, 63, 64, or 65, wherein said cytokine is interleukin.

67. The recombinant cell of claim 66, wherein said interleukin is 1L-2, 1L-4, or 1L-12.

68. The recombinant cell of claim 58, 59, 60, or 61, wherein said recombinant cell is a eukaryotic cell.

69. The recombinant cell of claim 68, which has been rendered non-proliferative.

70. The recombinant cell of claim 68, wherein said cell is a fibroblast.

71. Expression vector comprising a mutated or attenuated virus and the isolated nucleic acid molecule of claim 53, 54 or 55.

72. The expression vector of claim 71, wherein said virus is adenovirus, adeno associated virus, or vaccinia virus.

73. Expression system useful in making a recombinant cell, comprising:

(i) a first vector which encodes the protein encoded by the isolated nucleic acid molecule of claim 53, 54 or 55, and

(ii) a second vector which either (a) encodes an MHC or HLA molecule or (b) encodes an interleukin.

74. An isolated cancer associated antigen comprising the amino acid sequence encoded by SEQ ID NO: 5, 6, 7 or 8.

75. Composition comprising the isolated cancer associated antigen of claim 74, and a pharmaceutically acceptable adjuvant.

76. The composition of claim 75, wherein said adjuvant is a cytokine, a saponin, or GM-CSF.

77. Composition comprising at least one peptide consisting of an amino acid sequence of from 8 to 25 amino acids concatenated to each other in the isolated cancer associated antigen of claim 74, and a pharmaceutically acceptable adjuvant.

78. The composition of claim 77, wherein said adjuvant is a saponin, a cytokine, or GM-CSF.

79. The composition of claim 77, comprising a plurality of MHC binding peptides.

80. Composition comprising an expression vector which encodes at least one peptide consisting of an amino acid sequence of from 8 to 25 amino acids concatenated to each other in the isolated cancer associated antigen of claim 74, and pharmaceutically acceptable adjuvant.

81. The composition of claim 80, wherein said expression vector encodes a plurality of peptides.

82. Composition useful in treating a subject afflicted with a cancer, comprising the recombinant cell of claim 69 and a pharmacologically acceptable adjuvant.

83. The composition of claim 82, wherein said recombinant cell expresses an HLA or MHC molecule.

84. The composition of claim 82, wherein said recombinant cell is a human cell.

85. The composition of claim 77, further comprising at least one MHC or HLA molecule.

86. Isolated antibody which specifically binds to the isolated cancer associated antigen of claim 74.

87. The isolated antibody of claim 86, wherein said antibody is a monoclonal antibody.

88. A method for screening for possible presence of a pathological condition, comprising assaying a sample from a patient believed to have a pathological condition for antibodies specific to at least one of the cancer associated antigens encoded by SEQ ID NOS: 4, 5, 6, 7 or 8, presence of said antibodies being indicative of possible presence of said pathological condition.

89. The method of claim 88, wherein said pathological condition is cancer.

90. The method of claim 89, wherein said cancer is melanoma.

91. The method of claim 90, further comprising contacting said sample to purified cancer associated antigen encoded by SEQ ID NO: 4, 5, 6, 7 or 8.

92. A method for screening for possible presence of a pathological condition in a subject, comprising assaying a sample taken from said subject for expression of a nucleic acid molecule, the nucleotide sequence of which comprises SEQ ID NO: 5, 6, 7 or 8, expression of said nucleic acid molecule being indicative of possible presence of said pathological condition.

93. The method of claim 92, wherein said pathological condition is cancer.

94. The method of claim 92, comprising determining expression via polymerase chain reaction.

95. The method of claim 92, comprising determining expression by contacting said sample with at least one of SEQ ID NO: 11, 12, 13 or 14.



96. A method for determining regression, progression or onset of a cancerous condition comprising monitoring a sample from a patient with said cancerous condition for a parameter selected from the group consisting of (i) a cancer associated antigen encoded by SEQ ID NO: 3, 4, 5 or 6, (ii) a peptide derived from said cancer associated antigen, (iii) cytolytic T cells specific for said peptide and an MHC molecule with which it non-covalently complexes, and (iv) antibodies specific for said cancer associated antigen, wherein amount of said parameter is indicative of progression or regression or onset of said cancerous condition.

97. The method of claim 96, wherein said sample is a body fluid or exudate.

98. The method of claim 96, wherein said sample is a tissue.

99. The method of claim 96, comprising contacting said sample with an antibody which specifically binds with said protein or peptide.

100. The method of claim 99, wherein said antibody is labelled with a radioactive label or an enzyme.

101. The method of claim 99, wherein said antibody is a monoclonal antibody.

102. The method of claim 96, comprising amplifying RNA which codes for said protein.

103. The method of claim 102, wherein said amplifying comprises carrying out polymerase chain reaction.

104. The method of claim 96, comprising contacting said sample with a nucleic acid molecule which specifically hybridizes to a nucleic acid molecule which codes for or expresses said protein.

105. The method of claim 96, comprising assaying said sample for shed cancer associated antigen.

106. The method of claim 96, comprising assaying said sample for antibodies specific for said cancer associated antigen, by contacting said sample with said cancer associated antigen.

107. Method for screening for a cancerous condition comprising assaying a sample taken from a subject for an immunoreactive cell specific for a peptide derived from a cancer associated antigen encoded by SEQ ID NO: 4, 5, 6, 7 or 8 complexed to an MHC molecule, presence of said immunoreactive cell being indicative of said cancerous condition.

108. An isolated nucleic acid molecule consisting of a nucleotide sequence defined by SEQ ID NO: 9, 10, 11, 12, 13 or 14.

109. Kit useful in determining expression of a cancer associated antigen, comprising a separate portion of each of (i) the nucleotide sequences defined by SEQ ID NOS: 9 and 10, (ii) the nucleotide sequences defined by SEQ ID NOS: 11 and 12, and (iii) the nucleotide sequences defined by SEQ ID NOS: 13 and 14.

<110> Chen, Yao-Tseng  
Gure, Ali  
Tsang, Solam  
Stockert, Elisabeth  
Jager, Elke  
Knuth, Alexander  
Old, Lloyd J.

<120> Isolated Nucleic Acid Molecules Encoding Cancer Associated Antigen, The Antigens Per Se, And Uses Thereof

<130> LUD 5538.1 PCT

<140> PCT/US 99/05766

<141> 1998 - 04 - 17

<160> 3

<210> 1

<211> 4265

<212> DNA

<213> Homo sapiens

<220>

<400> 1

GTCTGAAGGA	CCTGAGGCAT	TTTGTGACGA	GGATCGTCTC	AGGTCAGCGG	AGGGAGGAGA	60
CTTATAGACC	TATCCAGTCT	TCAAGGTGCT	CCAGAAAGCA	GGAGTTGAAG	ACCTGGGTGT	120
GAGGGACACA	TACATCCTAA	AAGCACCACA	GCAGAGGAGG	CCCAGGCAGT	GCCAGGAGTC	180
AAGGTTCCCA	GAAGACAAAC	CCCCTAGGAA	GACAGGCGAC	CTGTGAGGCC	CTAGAGCACC	240
ACCTTAAGAG	AAGAAGAGCT	GTAAGCCGGC	CTTTGTCAGA	GCCATCATGG	GGGACAAGGA	300
TATGCCTACT	GCTGGGATGC	CGAGTCTTCT	CCAGAGTTCC	TCTGAGAGTC	CTCAGAGTTG	360
TCCTGAGGGG	GAGGACTCCC	AGTCTCCTCT	CCAGATTCCC	CAGAGTTCTC	CTGAGAGCGA	420
CGACACCCTG	TATCCTCTCC	AGAGTCCTCA	GAGTCGTTCT	GAGGGGGAGG	ACTCCTCGGA	480
TCCTCTCCAG	AGACCTCCTG	AGGGGAAGGA	CTCCCAGTCT	CCTCTCCAGA	TTCCCCAGAG	540
TTCTCCTGAG	GGCGACGACA	CCCAGTCTCC	TCTCCAGAAT	TCTCAGAGTT	CTCCTGAGGG	600
GAAGGACTCC	CTGTCTCCTC	TAGAGATTTT	TCAGAGCCCT	CCTGAGGGTG	AGGATGTCCA	660
GTCTCCTCTG	CAGAATCCTG	CGAGTTCCCT	CTTCTCCTCT	GCTTTATTGA	GTATTTTCCA	720
GAGTTCCCTT	GAGAGTATTC	AAAGTCCTTT	TGAGGGTTTT	CCCCAGTCTG	TTCTCCAGAT	780
TCCTGTGAGC	GCCGCCTCCT	CCTCCACTTT	AGTGAGTATT	TTCCAGAGTT	CCCCTGAGAG	840
TACTCAAAGT	CCTTTTGAGG	GTTTTCCCCA	GTCTCCACTC	CAGATTCTCT	TGAGCCGCTC	900
CTTCTCCTCC	ACTTTATTGA	GTATTTTCCA	GAGTTCCCTT	GAGAGAAGTC	AGAGAACTTC	960
TGAGGGTTTT	GCACAGTCTC	CTCTCCAGAT	TCCTGTGAGC	TCCTCCTCGT	CCTCCACTTT	1020
ACTGAGTCTT	TTCCAGAGTT	CCCCTGAGAG	AACTCAGAGT	ACTTTTGAGG	GTTTTCCCCA	1080
GTCTCCACTC	CAGATTCTCT	TGAGCCGCTC	CTTCTCCTCC	ACTTTATTGA	GTATTTTCCA	1140
GAGTTCCCTT	GAGAGAACTC	AGAGTACTTT	TGAGGGTTTT	GCCCAGTCTC	CTCTCCAGAT	1200
TCCTGTGAGC	CCCTCCTTCT	CCTCCACTTT	AGTGAGTATT	TTCCAGAGTT	CCCCTGAGAG	1260
AACTCAGAGT	ACTTTTGAGG	GTTTTCCCCA	GTCTCCTCTC	CAGATTCTCT	TGAGCTCCTC	1320
CTTCTCCTCC	ACTTTATTGA	GTCTTTTCCA	GAGTTCCCTT	GAGAGAACTC	AGAGTACTTT	1380
TGAGGGTTTT	CCCCAGTCTC	CTCTCCAGAT	TCCTGGAAGC	CCCTCCTTCT	CCTCCACTTT	1440
ACTGAGTCTT	TTCCAGAGTT	CCCCTGAGAG	AACTCACAGT	ACTTTTGAGG	GTTTTCCCCA	1500
GTCTCCTCTC	CAGATTCTCT	TGACCTCCTC	CTTCTCCTCT	ACTTTATTGA	GTATTTTACA	1560
GAGTTCTCCT	GAGAGTGCTC	AAAGTGCTTT	TGAGGGTTTT	CCCCAGTCTC	CTCTCCAGAT	1620
TCCTGTGAGC	TCCTCTTTCT	CCTACACTTT	ATTGAGTCTT	TTCCAGAGTT	CCCCTGAGAG	1680
AACTCACAGT	ACTTTTGAGG	GTTTTCCCCA	GTCTCCTCTC	CAGATTCTCT	TGAGCTCCTC	1740
CTCCTCCTCC	TCCACTTTAT	TGAGTCTTTT	CCAGAGTTCC	CCTGAGTGTA	CTCAAAGTAC	1800
TTTTGAGGGT	TTTCCCCAGT	CTCCTCTCCA	GATTCTCTCAG	AGTCCTCCTG	AAGGGGAGAA	1860
TACCCATTCT	CCTCTCCAGA	TTGTTCCAAG	TCTTCCTGAG	TGGGAGGACT	CCCTGTCTCC	1920
TCACTACTTT	CCTCAGAGCC	CTCCTCAGGG	GGAGGACTCC	CTATCTCCTC	ACTACTTTCC	1980
TCAGAGCCCT	CCTCAGGGGG	AGGACTCCCT	GTCTCCTCAC	TACTTTCCTC	AGAGCCCTCA	2040
GGGGGAGGAC	TCCCTGTCTC	CTCACTACTT	TCCTCAGAGC	CCTCCTCAGG	GGGAGGACTC	2100
CATGTCTCCT	CTCTACTTTC	CTCAGAGTCC	TCTTCAGGGG	GAGGAATTCC	AGTCTTCTCT	2160
CCAGAGCCCT	GTGAGCATCT	GCTCCTCCTC	CACTCCATCC	AGTCTTCCCC	AGAGTTTCCC	2220
TGAGAGTTCT	CAGAGTCCTC	CTGAGGGGCC	TGTCCAGTCT	CCTCTCCATA	GTCTCAGAG	2280

```

CCCTCCTGAG GGGATGCACT CCCAATCTCC TCTCCAGAGT CCTGAGAGTG CTCCTGAGGG 2340
GGAGGATTCC CTGTCTCCTC TCCAAATTCC TCAGAGTCCT CTTGAGGGAG AGGACTCCCT 2400
GTCTTCTCTC CATTTTCTCTC AGAGTCCTCC TGAGTGGGAG GACTCCCTCT CTCCTCTCCA 2460
CTTTCCTCAG TTTCTCCTCTC AGGGGGAGGA CTTCCAGTCT TCTCTCCAGA GTCCTGTGAG 2520
TATCTGCTCC TCCTCCACTT CTTTGAGTCT TCCCCAGAGT TTCCCTGAGA GTCCTCAGAG 2580
TCCTCCTGAG GGGCCTGCTC AGTCTCCTCT CCAGAGACCT GTCAGCTCCT TCTTCTCCTA 2640
CACTTTAGCG AGTCTTCTCC AAAGTTCCCA TGAGAGTCCT CAGAGTCCTC CTGAGGGGCC 2700
TGCCCACTCT CCTCTCCAGA GTCCTGTGAG CTCCTTCCCC TCCTCCACTT CATCGAGTCT 2760
TTCCCAGAGT TCTCCTGTGA GCTCCTTCCC CTCCTCCACT TCATCGAGTC TTTCCAAGAG 2820
TTCCCCTGAG AGTCCTCTCC AGAGTCCTGT GATCTCCTTC TCCTCCTCCA CTTTATTGAG 2880
CCCATTCACT GAAGAGTCCA GCAGCCCACT AGATGAATAT ACAAGTTCCT CAGACACCTT 2940
GCTAGAGAGT GATTCTCTGA CAGACAGCGA GTCCTTGATA GAGAGCGAGC CCTTGTTTAC 3000
TTATACACTG GATGAAAAGG TGGACGAGTT GGC GCGGTTT CTTCTCCTCA AATATCAAGT 3060
GAAGCAGCCT ATCACAAGG CAGAGATGCT GACGAATGTC ATCAGCAGGT ACACGGGCTA 3120
CTTTCCTGTG ATCTTCAGGA AAGCCCGTGA GTTCATAGAG ATACTTTTTG GCATTTCCCT 3180
GAGAGAAGTG GACCCTGATG ACTCCTATGT CTTTGTAAC ACATTAGACC TCACCTCTGA 3240
GGGGTGCTG AGTGATGAGC AGGGCATGTC CCAGAACCGC CTCCTGATTC TTATTCTGAG 3300
TATCATCTTC ATAAAGGGCA CCTATGCCTC TGAGGAGGTC ATCTGGGATG TGCTGAGTGG 3360
AATAGGGGTG CGTGCTGGGA GGGAGCACTT TGCCTTTGGG GAGCCAGGG AGCTCCTCAC 3420
TAAAGTTTGG GTGCAGGAAC ATTACCTAGA GTACCGGGAG GTGCCAACT CTTCTCCTCC 3480
TCGTTACGAA TTCCTGTGGG GTCCAAGAGC TCATTACAGAA GTCATTAAGA GGAAAGTAGT 3540
AGAGTTTTTG GCCATGCTAA AGAATACCGT CCCTATTACC TTTCCATCCT CTTACAAGGA 3600
TGCTTTGAAA GATGTGGAAG AGAGAGCCCA GGCCATAATT GACACCACAG ATGATTCTGAC 3660
TGCCACAGAA AGTGCAAGCT CCAGTGTCAT GTCCCCCAGC TTCTCTTCTG AGTGAACTCT 3720
AGGGCAGATT CTTCCCTCTG AGTTTGAAGG GGGCAGTCGA GTTTCTACGT GGTGGAGGGC 3780
CTGGTTGAGG CTGGAGAGAA CACAGTGCTA TTTGCATTTT TGTTCCATAT GGGTAGTTAT 3840
GGGGTTTACC TGTTTTACTT TTGGGTATTT TTCAAATGCT TTTCTTATTA ATAACAGGTT 3900
TAAATAGCTT CAGAATCCTA GTTTATGCAC ATGAGTCGCA CATGTATTGC TGTTTTCTG 3960
GTTTAAGAGT AACAGTTTGA TATTTTGTAA AAACAAAAC ACACCCAAAC ACACCACATT 4020
GGGAAAACCT TCTGCCTCAT TTTGTGATGT GTACAGGTT AATGTGGTGT TACTGTAGGA 4080
ATTTTCTTGA AACTGTGAAG GAACTCTGCA GTTAAATAGT GGAATAAAGT AAAGGATTGT 4140
TAATGTTTGC ATTTCTCAG GTCCTTTAGT CTGTTGTTCT TGAAACTAA AGATACATAC 4200
CTGGTTTGCT TGGCTTACGT AAGAAAGTCG AAGAAAGTAA ACTGTAATAA ATAAAAGTGT 4260
CAGTG 4265

```

```

<210>      2
<211>      1142
<212>      PRT
<213>      Homo sapiens
<220>
<400>      2

```

```

Met Gly Asp Lys Asp Met Pro Thr Ala Gly Met Pro Ser Leu Leu Gln
                    5                      10                      15
Ser Ser Ser Glu Ser Pro Gln Ser Cys Pro Glu Gly Glu Asp Ser Gln
                    20                      25                      30
Ser Pro Leu Gln Ile Pro Gln Ser Ser Pro Glu Ser Asp Thr Leu
                    35                      40                      45
Tyr Pro Leu Gln Ser Pro Gln Ser Arg Ser Glu Gly Glu Asp Ser Ser
                    50                      55                      60
Asp Pro Leu Gln Arg Pro Pro Glu Gly Lys Asp Ser Gln Ser Pro Leu
                    65                      70                      75                      80
Gln Ile Pro Gln Ser Ser Pro Glu Gly Asp Thr Gln Ser Pro Leu
                    85                      90                      95
Gln Asn Ser Gln Ser Ser Pro Glu Gly Lys Asp Ser Leu Ser Pro Leu
                    100                      105                      110
Glu Ile Ser Gln Ser Pro Pro Glu Gly Glu Asp Val Gln Ser Pro Leu
                    115                      120                      125
Gln Asn Pro Ala Ser Ser Phe Phe Ser Ser Ala Leu Leu Ser Ile Phe
                    130                      135                      140
Gln Ser Ser Pro Glu Ser Ile Gln Ser Pro Phe Glu Gly Phe Pro Gln
                    145                      150                      155                      160
Ser Val Leu Gln Ile Pro Val Ser Ala Ala Ser Ser Ser Thr Leu Val
                    165                      170                      175

```

## SUBSTITUTE SHEET (Rule 26)

Ser Ile Phe Gln Ser Ser Pro Glu Ser Thr Gln Ser Pro Phe Glu Gly  
 180 185 190  
 Phe Pro Gln Ser Pro Leu Gln Ile Pro Val Ser Arg Ser Phe Ser Ser  
 195 200 205  
 Thr Leu Leu Ser Ile Phe Gln Ser Ser Pro Glu Arg Ser Gln Arg Thr  
 210 215 220  
 Ser Glu Gly Phe Ala Gln Ser Pro Leu Gln Ile Pro Val Ser Ser Ser  
 225 230 235 240  
 Ser Ser Ser Thr Leu Leu Ser Leu Phe Gln Ser Ser Pro Glu Arg Thr  
 245 250 255  
 Gln Ser Thr Phe Glu Gly Phe Pro Gln Ser Pro Leu Gln Ile Pro Val  
 260 265 270  
 Ser Arg Ser Phe Ser Ser Thr Leu Leu Ser Ile Phe Gln Ser Ser Pro  
 275 280 285  
 Glu Arg Thr Gln Ser Thr Phe Glu Gly Phe Ala Gln Ser Pro Leu Gln  
 290 295 300  
 Ile Pro Val Ser Pro Ser Phe Ser Ser Thr Leu Val Ser Ile Phe Gln  
 305 310 315 320  
 Ser Ser Pro Glu Arg Thr Gln Ser Thr Phe Glu Gly Phe Pro Gln Ser  
 325 330 335  
 Pro Leu Gln Ile Pro Val Ser Ser Ser Phe Ser Ser Thr Leu Leu Ser  
 340 345 350  
 Leu Phe Gln Ser Ser Pro Glu Arg Thr Gln Ser Thr Phe Glu Gly Phe  
 355 360 365  
 Pro Gln Ser Pro Leu Gln Ile Pro Gly Ser Pro Ser Phe Ser Ser Thr  
 370 375 380  
 Leu Leu Ser Leu Phe Gln Ser Ser Pro Glu Arg Thr His Ser Thr Phe  
 385 390 395 400  
 Glu Gly Phe Pro Gln Ser Pro Leu Gln Ile Pro Met Thr Ser Ser Phe  
 405 410 415  
 Ser Ser Thr Leu Leu Ser Ile Leu Gln Ser Ser Pro Glu Ser Ala Gln  
 420 425 430  
 Ser Ala Phe Glu Gly Phe Pro Gln Ser Pro Leu Gln Ile Pro Val Ser  
 435 440 445  
 Ser Ser Phe Ser Tyr Thr Leu Ser Leu Phe Gln Ser Ser Pro Glu  
 450 455 460  
 Arg Thr His Ser Thr Phe Glu Gly Phe Pro Gln Ser Pro Leu Gln Ile  
 465 470 475 480  
 Pro Val Ser Ser Ser Ser Ser Ser Ser Thr Leu Leu Ser Leu Phe Gln  
 485 490 495  
 Ser Ser Pro Glu Cys Thr Gln Ser Thr Phe Glu Gly Phe Pro Gln Ser  
 500 505 510  
 Pro Leu Gln Ile Pro Gln Ser Pro Pro Glu Gly Glu Asn Thr His Ser  
 515 520 525  
 Pro Leu Gln Ile Val Pro Ser Leu Pro Glu Trp Glu Asp Ser Leu Ser  
 530 535 540  
 Pro His Tyr Phe Pro Gln Ser Pro Pro Gln Gly Glu Asp Ser Leu Ser  
 545 550 555 560  
 Pro His Tyr Phe Pro Gln Ser Pro Pro Gln Gly Glu Asp Ser Leu Ser  
 565 570 575  
 Pro His Tyr Phe Pro Gln Ser Pro Gln Gly Glu Asp Ser Leu Ser Pro  
 580 585 590  
 His Tyr Phe Pro Gln Ser Pro Pro Gln Gly Glu Asp Ser Met Ser Pro  
 595 600 605  
 Leu Tyr Phe Pro Gln Ser Pro Leu Gln Gly Glu Glu Phe Gln Ser Ser  
 610 615 620  
 Leu Gln Ser Pro Val Ser Ile Cys Ser Ser Ser Thr Pro Ser Ser Leu  
 625 630 635 640  
 Pro Gln Ser Phe Pro Glu Ser Ser Gln Ser Pro Pro Glu Gly Pro Val  
 645 650 655  
 Gln Ser Pro Leu His Ser Pro Gln Ser Pro Pro Glu Gly Met His Ser  
 660 665 670  
 Gln Ser Pro Leu Gln Ser Pro Glu Ser Ala Pro Glu Gly Glu Asp Ser  
 675 680 685

Leu Ser Pro Leu Gln Ile Pro Gln Ser Pro Leu Glu Gly Glu Asp Ser  
 690 695 700  
 Leu Ser Ser Leu His Phe Pro Gln Ser Pro Pro Glu Trp Glu Asp Ser  
 705 710 715  
 Leu Ser Pro Leu His Phe Pro Gln Phe Pro Pro Gln Gly Glu Asp Phe  
 725 730 735  
 Gln Ser Ser Leu Gln Ser Pro Val Ser Ile Cys Ser Ser Ser Thr Ser  
 740 745 750  
 Leu Ser Leu Pro Gln Ser Phe Pro Glu Ser Pro Gln Ser Pro Pro Glu  
 755 760 765  
 Gly Pro Ala Gln Ser Pro Leu Gln Arg Pro Val Ser Ser Phe Phe Ser  
 770 775 780  
 Tyr Thr Leu Ala Ser Leu Leu Gln Ser Ser His Glu Ser Pro Gln Ser  
 785 790 795 800  
 Pro Pro Glu Gly Pro Ala Gln Ser Pro Leu Gln Ser Pro Val Ser Ser  
 805 810 815  
 Phe Pro Ser Ser Thr Ser Ser Ser Leu Ser Gln Ser Ser Pro Val Ser  
 820 825 830  
 Ser Phe Pro Ser Ser Thr Ser Ser Ser Leu Ser Lys Ser Ser Pro Glu  
 835 840 845  
 Ser Pro Leu Gln Ser Pro Val Ile Ser Phe Ser Ser Ser Thr Ser Leu  
 850 855 860  
 Ser Pro Phe Ser Glu Glu Ser Ser Ser Pro Val Asp Glu Tyr Thr Ser  
 865 870 875 880  
 Ser Ser Asp Thr Leu Leu Glu Ser Asp Ser Leu Thr Asp Ser Glu Ser  
 885 890 895  
 Leu Ile Glu Ser Glu Pro Leu Phe Thr Tyr Thr Leu Asp Glu Lys Val  
 900 905 910  
 Asp Glu Leu Ala Arg Phe Leu Leu Lys Tyr Gln Val Lys Gln Pro  
 915 920 925  
 Ile Thr Lys Ala Glu Met Leu Thr Asn Val Ile Ser Arg Tyr Thr Gly  
 930 935 940  
 Tyr Phe Pro Val Ile Phe Arg Lys Ala Arg Glu Phe Ile Glu Ile Leu  
 945 950 955 960  
 Phe Gly Ile Ser Leu Arg Glu Val Asp Pro Asp Asp Ser Tyr Val Phe  
 965 970 975  
 Val Asn Thr Leu Asp Leu Thr Ser Glu Gly Cys Leu Ser Asp Glu Gln  
 980 985 990  
 Gly Met Ser Gln Asn Arg Leu Leu Ile Leu Ile Leu Ser Ile Ile Phe  
 995 1000 1005  
 Ile Lys Gly Thr Tyr Ala Ser Glu Glu Val Ile Trp Asp Val Leu Ser  
 1010 1015 1020  
 Gly Ile Gly Val Arg Ala Gly Arg Glu His Phe Ala Phe Gly Glu Pro  
 1025 1030 1035 1040  
 Arg Glu Leu Leu Thr Lys Val Trp Val Gln Glu His Tyr Leu Glu Tyr  
 1045 1050 1055  
 Arg Glu Val Pro Asn Ser Ser Pro Pro Arg Tyr Glu Phe Leu Trp Gly  
 1060 1065 1070  
 Pro Arg Ala His Ser Glu Val Ile Lys Arg Lys Val Val Glu Phe Leu  
 1075 1080 1085  
 Ala Met Leu Lys Asn Thr Val Pro Ile Thr Phe Pro Ser Ser Tyr Lys  
 1090 1095 1100  
 Asp Ala Leu Lys Asp Val Glu Glu Arg Ala Gln Ala Ile Ile Asp Thr  
 1105 1110 1115 1120  
 Thr Asp Asp Ser Thr Ala Thr Glu Ser Ala Ser Ser Ser Val Met Ser  
 1125 1130 1135  
 Pro Ser Phe Ser Ser Glu  
 1140

<210> 3  
 <211> 7  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <400> 3

Pro Gln Ser Pro Leu Gln Ile  
 1 5

<210> 4  
 <211> 4159  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <400> 4

GGTGGATGCG	TTTGGGTTGT	AGCTAGGCTT	TTTCTTTTCT	TTCTCTTTTA	AAACACATCT	60
AGACAAAGGAA	AAAACAAGCC	TCGGATCTGA	TTTTTCACTC	CTCGTTCTTG	TGCTTGGTTC	120
TTACTGTGTT	TGTGTATTTT	AAAGGCGAGA	AGACGAGGGG	AACAAAACCA	GCTGGATCCA	180
TCCATCACCG	TGGGTGGTTT	TAATTTTTCG	TTTTTTCTCG	TTATTTTTTT	TTAAACAACC	240
ACTCTTCACA	ATGAACAAAC	TGTATATCGG	AAACCTCAGC	GAGAACGCCG	CCCCCTCGGA	300
CCTAGAAAGT	ATCTTCAAGG	ACGCCAAGAT	CCCGGTGTCG	GGACCCTTCC	TGGTGAAGAC	360
TGGCTACGCG	TTCGTGGACT	GCCCGGACGA	GAGCTGGGCC	CTCAAGGCCA	TCGAGGCGCT	420
TTCAGGTAAA	ATAGAACTGC	ACGGGAAACC	CATAGAAGTT	GAGCACTCGG	TCCCAAAAAG	480
GCAAAGGATT	CGGAAACTTC	AGATACGAAA	TATCCCGCCT	CATTTACAGT	GGGAGGTGCT	540
GGATAGTTTA	CTAGTCCAGT	ATGGAGTGGT	GGAGAGCTGT	GAGCAAGTGA	ACACTGACTC	600
GGAAACTGCA	GTTGTAAATG	TAACCTATTC	CAGTAAGGAC	CAAGCTAGAC	AAGCACTAGA	660
CAAACCTGAAT	GGATTTTCAGT	TAGAGAATTT	CACCTTGAAA	GTAGCCTATA	TCCCTGATGA	720
AATGGCCGCC	CAGCAAAACC	CCTTGCAGCA	GCCCCGAGGT	CGCCGGGGGC	TTGGGCAGAG	780
GGGCTCCTCA	AGGCAGGGGT	CTCCAGGATC	CGTATCCAAG	CAGAAACCAT	GTGATTTGCC	840
TCTGCGCCTG	CTGGTTCCCA	CCCAATTTGT	TGGAGCCATC	ATAGGAAAAG	AAGGTGCCAC	900
CATTTCGGAA	ATCACCAAAC	AGACCCAGTC	TAAAATCGAT	GTCCACCGTA	AAGAAAATGC	960
GGGGGCTGCT	GAGAAGTCGA	TTACTATCCT	CTCTACTCCT	GAAGGCACCT	CTGCGGCTTG	1020
TAAGTCTATT	CTGGAGATTA	TGCATAAGGA	AGCTCAAGAT	ATAAAATTCA	CAGAAGAGAT	1080
CCCCTTGAAG	ATTTTAGCTC	ATAATAACTT	TGTTGGACGT	CTTATTGGTA	AAGAAGGAAG	1140
AAATCTTAAA	AAAATTGAGC	AAGACACAGA	CACTAAAATC	ACGATATCTC	CATTGCAGGA	1200
ATTGACGCTG	TATAATCCAG	AACGCACTAT	TACAGTTAAA	GGCAATGTTG	AGACATGTGC	1260
CAAAGCTGAG	GAGGAGATCA	TGAAGAAAAT	CAGGGAGTCT	TATGAAAATG	ATATTGCTTC	1320
TATGAATCTT	CAAGCACATT	TAATTCCTGG	ATTAAATCTG	AACGCCTTGG	GTCTGTTCCC	1380
ACCCACTTCA	GGGATGCCAC	CTCCCACCTC	AGGGCCCCCT	TCAGCCATGA	CTCCTCCCTA	1440
CCCGCAGTTT	GAGCAATCAG	AAACGGGAGC	TGTTTCATCAG	TTTATCCCAG	CTCTATCAGT	1500
CGGTGCCATC	ATCGGCAAGC	AGGGCCAGCA	CATCAAGCAG	CTTTCTCGCT	TTGCTGGAGC	1560
TTCAATTAAG	ATTGCTCCAG	CGGAAGCAC	AGATGCTAAA	GTGAGGATGG	TGATTATCAC	1620
TGGACCAACA	GAGGCTCAGT	TCAAGGCTCA	GGGAAGAATT	TATGGAAAAA	TTAAAGAAGA	1680
AAACTTTGTT	AGTCCTAAAG	AAGAGGTGAA	ACTTGAAGCT	CATATCAGAG	TGCCATCCTT	1740
TGCTGCTGGC	AGAGTTATTG	GAAAAGGAGG	CAAAACGGTG	AATGAACCTC	AGAATTTGTC	1800
AAGTGCAGAA	GTTGTTGTCC	CTCGTGACCA	GACACCTGAT	GAGAATGACC	AAGTGGTTGT	1860
CAAAATAACT	GGTCACTTCT	ATGCTTGCCA	GGTTGCCAG	AGAAAAATTC	AGGAAATTC	1920
GACTCAGGTA	AAGCAGCAC	AACAACAGAA	GGCTCTGCAA	AGTGGACCAC	CTCAGTCAAG	1980
ACGGAAGTAA	AGGCTCAGGA	AACAGCCCAC	CACAGAGGCA	GATGCCAAAC	CAAAGACAGA	2040
TTGCTTAACC	AACAGATGGG	CGCTGACCCC	CTATCCAGAA	TCACATGCAC	AAGTTTTTAC	2100
CTAGCCAGTT	GTTTCTGAGG	ACCAGGCAAC	TTTTGAACTC	CTGTCTCTGT	GAGAATGTAT	2160
ACTTTATGCT	CTCTGAAATG	TATGACACCC	AGCTTTAAAA	CAAACAAACA	AACAAACAAA	2220
AAAAGGGTGG	GGGAGGGAGG	GAAAGAGAAG	AGCTCTGCAC	TTCCCTTTGT	TGTAGTCTCA	2280
CAGTATAACA	GATATTCTAA	TTCTTCTTAA	GATTTCCCCA	TAATGCCAGA	AATTGGCTTA	2340
ATGATGCTTT	CACTAAATTC	ATCAAATAGA	TTGCTCCTAA	ATCCAATTGT	TAAAATTGGA	2400
TCAGAATAAT	TATCACAGGA	ACTTAAATGT	TAAGCCATTA	GCATAGAAAA	ACTGTTCTCA	2460
GTTTTATTTT	TACCTAACAC	TAACATGAGT	AACCTAAGGG	AAGTGCTGAA	TGGTGTTGGC	2520
AGGGGTATTA	AACGTGCATT	TTTACTCAAC	TACCTCAGGT	ATTGAGTAAT	ACAATGAAAA	2580
GCAAAATTGT	TCCTTTTTTT	TGAAAATTTT	ATATACTTTA	TAATGATAGA	AGTCCAACCG	2640
TTTTTTAAAA	AATAAATTTA	AAATTTAACA	GCAATCAGCT	AACAGGCAAA	TTAAGATTTT	2700

TACTTCTGGC	TGGTGACAGT	AAAGCTGGAA	AATTAATTTT	AGGGTTTTTT	GAGGCTTTTG	2760
ACACAGTTAT	TAGTTAAATC	AAATGTTCAA	AAATACGGAG	CAGTGCCTAG	TATCTGGAGA	2820
GCAGCACTAC	CATTTATTCT	TTCATTTATA	GTTGGGAAAG	TTTTTGACGG	TACTAACAAA	2880
GTGGTCGCAG	GAGATTTTGG	AACGGCTGGT	TTAAATGGCT	TCAGGAGACT	TCAGTTTTTT	2940
GTTTAGCTAC	ATGATTGAAT	GCATAATAAA	TGCTTTGTGC	TTCTGACTAT	CAATACCTAA	3000
AGAAAGTGCA	TCAGTGAAGA	GATGCAAGAC	TTTCAACTGA	CTGGCAAAAA	GCAAGCTTTA	3060
GCTTGTCTTA	TAGGATGCTT	AGTTTGCCAC	TACACTTCAG	ACCAATGGGA	CAGTCATAGA	3120
TGGTGTGACA	GTGTTTAAAC	GCAACAAAAG	GCTACATTTT	CATGGGGCCA	GCACTGTCAT	3180
GAGCCTCACT	AAGCTATTTT	GAAGATTTTT	AAGCACTGAT	AAATTAAAAA	AAAAAAAAAA	3240
AAATTAGACT	CCACCTTAAG	TAGTAAAGTA	TAACAGGATT	TCTGTATACT	GTGCAATCAG	3300
TTCTTTGAAA	AAAAAGTCAA	AAGATAGAGA	ATACAAGAAA	AGTTTNGGG	ATATAATTGG	3360
AATGACTGTG	AAAACATATG	ACCTTTGATA	ACGAACCTAT	TTGCTCACTC	CTTGACAGCA	3420
AAGCCCACTA	CGTACAATTG	TGTTGGGTGT	GGGTGGTCTC	CAAGGCCACG	CTGCTCTCTG	3480
AATTGATTTT	TTGAGTTTTG	GNTTGNAAGA	TGATCACAGN	CATGTTACAC	TGATCTTNAA	3540
GGACATATNT	TATAACCCTT	TAAAAAATAA	ATCCCCTGCC	TCATTCTTAT	TTCGAGATGA	3600
ATTTCGATAC	AGACTAGATG	TCTTTCTGAA	GATCAATTAG	ACATTNTGAA	AATGATTTAA	3660
AGTGTTTTCC	TTAATGTTCT	CTGAAAACAA	GTTTCTTTTG	TAGTTTTAAC	CAAAAAAGTG	3720
CCCTTTTTGT	CACGTGGTTT	TCCTAGCATT	CATGATTTT	TTTCACACA	ATGAATTAAA	3780
ATTGCTAAAA	TCATGGACTG	GCTTTCTGGT	TGGATTTTCT	GTAAGATGTG	TTTAAGGCCA	3840
GAGCTTTTCT	CAGTATTTGA	TTTTTTTCCC	CAATATTTGA	TTTTTTAAAA	ATATACACAT	3900
AGGAGCTGCA	TTTAAACCTT	GCTGGTTTTA	ATTCTGTGAN	ATTTCACTTC	TAGCCTTTTA	3960
GTATGGCNAA	TCANAATTTA	CTTTTACTTA	AGCATTGTGA	ATTTGGAGTA	TCTGGTACTA	4020
GCTAAGAAAT	AATTGNATAA	TTGAGTTTTG	TACTCNCCAA	ANATGGGTCA	TTCTCATGN	4080
ATAATGTNCC	CCCAATGCAG	CTTCATTTTC	CAGANACCTT	GACGCAGGAT	AAATTTTTTC	4140
ATCATTTAGG	TCCCCAAA					4159

<210> 5  
 <211> 1708  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <400> 5

AGGGACGCTG	CCGCACCGCC	CCAGTTTACC	CCGGGGAGCC	ATCATGAAGC	TGAATGGCCA	60
CCAGTTGGAG	AACCATGCC	TGAAGGTCTC	CTACATCCCC	GATGAGCAGA	TAGCACAGGG	120
ACCTGAGAAT	GGGCGCCGAG	GGGGCTTTGG	CTCTCGGGGT	CAGCCCCGCC	AGGGCTCACC	180
TGTGGCAGCG	GGGGCCCCAG	CCAAGCAGCA	GCAAGTGGAC	ATCCCCCTTC	GGCTCCTGGT	240
GCCCACCCAG	TATGTGGGTG	CCATTATTGG	CAAGGAGGGG	GCCACCATCC	GCAACATCAC	300
AAAACAGACC	CAGTCCAAGA	TAGACGTGCA	TAGGAAGGAG	AACGCAGGTG	CAGCTGAAAA	360
AGCCATCAGT	GTGCACTCCA	CCCCTGAGGG	CTGCTCCTCC	GCTTGTAAGA	TGATCTTGGG	420
GATTATGCAT	AAAGAGGCTA	AGGACACCAA	AACGGCTGAC	GAGGTTCCCC	TGAAGATCCT	480
GGCCCATAA	AACTTTGTAG	GGCGTCTCAT	TGGCAAGGAA	GGACGGAACC	TGAAGAAGGT	540
AGAGCAAGAT	ACCGAGACAA	AAATCACCAT	CTCCTCGTTG	CAAGACCTTA	CCCTTTACAA	600
CCCTGAGAGG	ACCATCACTG	TGAAGGGGGC	CATCGAGAAT	TGTTGCAGGG	CCGAGCAGGA	660
AATAATGAAG	AAAGTTCGGG	AGGCCTATGA	GAATGATGTG	GCTGCCATGA	GCTCTCACCT	720
GATCCCTGGC	CTGAACCTGG	CTGCTGTAGG	TCTTTTCCCA	GCTTCATCCA	GCGCAGTCCC	780
GCCGCTCCC	AGCAGCGTTA	CTGGGGCTGC	TCCCTATAGC	TCCTTTATGC	AGGCTCCCGA	840
GCAGGAGATG	GTGCAGGTGT	TTATCCCCGC	CCAGGCAGTG	GGCGCCATCA	TCGCAAGAA	900
GGGGCAGCAC	ATCAAACAGC	TCTCCCGGTT	TGCCAGCGCC	TCCATCAAGA	TTGCACCACC	960
CGAAACACCT	GACTCCAAAG	TTCGTATGGT	TATCATCACT	GGACCGCCAG	AGGCCCAATT	1020
CAAGGCTCAG	GGAAGAATCT	ATGGCAAAC	CAAGGAGGAG	AACTTCTTTG	GTCCCAAGGA	1080
GGAAGTGAAG	CTGGAGACCC	ACATACGTGT	GCCAGCATCA	GCAGCTGGCC	GGGTCATTGG	1140
CAAGAGTGGA	AAAACGGTGA	ACGAGTTGCA	GCAATTTGACG	GCAGCTGAGG	TGGTAGTACC	1200
AAGAGACCAG	ACCCCTGATG	AGAACGACCA	GGTCATCGTG	AAAATCATCG	GACATTTCTA	1260
TGCCAGTCAG	ATGGCTCAAC	GGAAGATCCG	AGACATCCTG	GCCAGGTTA	AGCAGCAGCA	1320
TCAGAAGGGA	CAGAGTAACC	AGGCCAGGC	ACGGAGGAAG	TGACCAGCCC	CTCCCTGTCC	1380
CTTNGAGTCC	AGGACAACAA	CGGGCAGAAA	TCGAGAGTGT	GCTCTCCCCG	GCAGGCCTGA	1440
GAATGAGTGG	GAATCCGGGA	CACNTGGGCC	GGGCTGTAGA	TCAGGTTTGC	CCACTTGATT	1500
GAGAAAGATG	TTCCAGTGAG	GAACCTGTAT	CTNTCAGCCC	CAAACACCCA	CCCAATTGGC	1560
CCAACACTGT	NTGCCCTCG	GGGTGTCAGA	AATTNTAGCG	CAAGGCACTT	TAAACGTGG	1620
ATTGTTTAAA	GAAGCTCTCC	AGGCCCCACC	AAGAGGGTGG	ATCACACCTC	AGTGGGAAGA	1680
AAAATAAAAT	TTCCTTCAGG	TTTTAAAA				1708



<210> 6  
 <211> 3412  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <400> 6

```

GGCAGCGGAG GAGGCGAGGA GCGCCGGGTA CCGGGCCGGG GGAGCCGCGG GCTCTCGGGG 60
AAGAGACGGA TGATGAACAA GCTTTACATC GGGAACCTGA GCCCCGCCGT CACCGCCGAC 120
GACCTCCGGC AGCTCTTTGG GGACAGGAAG CTGCCCCCTG GGGACAGGT CCTGCTGAAG 180
TCCGGCTACG CCTTCGTGGA CTACCCCGAC CAGAACTGGG CCATCCGCGC CATCGAGACC 240
CTCTCGGGTA AAGTGGAAAT GCATGGGAAA ATCATGGAAG TTGATTACTC AGTCTCTAAA 300
AAGCTAAGGA GCAGGAAAAT TCAGATTCTGA AACATCCCTC CTCACCTGCA GTGGGAGGTG 360
TTGGATGGAC TTTTGGCTCA ATATGGGACA GTGGAGAATG TGGAACAAGT CAACACAGAC 420
ACAGAAACCG CCGTTGTCAA CGTCACATAT GCAACAAGAG AAGAAGCAAA AATAGCCATG 480
GAGAAGCTAA GCGGGCATCA GTTTGAGAAC TACTCCTTCA AGATTTCTTA CATCCCGGAT 540
GAAGAGGTGA GCTCCCCCTC GCCCCCTCAG CGAGCCCAGC GTGGGGACCA CTCTCCCGG 600
GAGCAAGGCC ACGCCCCTGG GGGCACTTCT CAGGCCAGAC AGATTGATTT CCCGCTGCGG 660
ATCCTGGTCC CCACCCAGTT TGTGTTGCC ATCATCGGAA AGGAGGGCTT GACCATAAAG 720
AACATCACTA AGCAGACCCA GTCCCGGGTA GATATCCATA GAAAAGAGAA CTCTGGAGCT 780
GCAGAGAAGC CTGTACCATT CCATGCCACC CCAGAGGGGA CTTCTGAAGC ATGCCGCATG 840
ATTCTTAAAA TCATGCAGAA AGAGGCAGAT GAGACCAAA TAGCCGAAGA GATTCCTCTG 900
AAAATCTTGG CACACAATGG CTTGGTTGGA AGACTGATTG GAAAAGAAGG CAGAAATTTG 960
AAGAAAATTG AACATGAAAC AGGGACCAAG ATAACAATCT CATCTTTGCA GGATTTGAGC 1020
ATATACAACC CGGAAAGAAC CATCACTGTG AAGGGCACAG TTGAGGCCTG TGCCAGTGCT 1080
GAGATAGAGA TTATGAAGAA GCTGCGTGAG GCCTTTGAAA ATGATATGCT GGCTGTAAAC 1140
CAACAAGCCA ATCTGATCCC AGGGTTGAAC CTCAGCGCAC TTGGCATCTT TCAACAGGA 1200
CTGTCCGTGC TATCTCCACC AGCAGGGCCC CGCGGAGCTC CCCCCGCTGC CCCCTACCAC 1260
CCCTTCACTA CCCACTCCGG ATACTTCTCC AGCCTGTACC CCCATCACC GTTTGGCCCG 1320
TTCCCGCATC ATCACTCTTA TCCAGAGCAG GAGATTGTGA ATCTCTTCAT CCCAACCCAG 1380
GCTGTGGGCG CCATCATCGG GAAGAAGGGG GCACACATCA AACAGCTGGC GAGATTGCGC 1440
GGAGCCTCTA TCAAGATTGC CCCTGCGGAA GGCCAGACG TCAGCGAAAG GATGGTCATC 1500
ATCACCGGGC CACCGGAAGC CCAGTTCAAG GCCCAGGGAC GGATCTTTGG GAAACTGAAA 1560
GAGGAAAAC TCTTTAACCC CAAAGAAGAA GTGAAGCTGG AAGCGCATAT CAGAGTGCCC 1620
TCTTCCACAG CTGGCCGGGT GATTGGCAAA GTTGGCAAGA CCGTGAACGA ACTGCAGAAC 1680
TTAACCAGTG CAGAAGTCAT CGTGCCCTCGT GACCAAACGC CAGATGAAAA TGAGGAAGTG 1740
ATCGTCAGAA TTATCGGGCA CTTCTTTGCT AGCCAGACTG CACAGCGCAA GATCAGGGAA 1800
ATTGTACAAC AGGTGAAGCA GCAGGAGCAG AAATACCTC AGGGAGTCGC CTCACAGCGC 1860
AGCAAGTGAG GCTCCACAG GCACCAGCAA AACAAACGGAT GAATGTAGCC CTTCCAACAG 1920
CTGACAGAAT GAGACCAAA GCAGCCAGCC AGATCGGGAG CAAACCAAAG ACCATCTGAG 1980
GAATGAGAAG TCTGCGGAGG CGGCCAGGGA CTCTGCCGAG GCCCTGAGAA CCCAGGGGC 2040
CGAGGAGGGG CGGGGAAGGT CAGCCAGGTT TGCCAGAACC ACCGAGCCCC GCCTCCCGCC 2100
CCCCAGGGCT TCTGCAGGCT TCAGCCATCC ACTTCACCAT CCACTCGGAT CTCTCCTGAA 2160
CTCCCACGAC GCTATCCCTT TTAGTTGAAC TAACATAGGT GAACGTGTTT AAAGCCAAGC 2220
AAAATGCACA CCCTTTTTCT GTGGCAAATC GTCTCTGTAC ATGTGTGTAC ATATTAGAAA 2280
GGGAAGATGT TAAGATATGT GGCCTGTGGG TTACACAGGG TGCCTGCAGC GGTAAATATAT 2340
TTTAGAAAA ATATATCAAA TAACTCAACT AACTCCAATT TTTAATCAAT TATTAATTTT 2400
TTTTTCTTTT TAAAGAGAAA GCAGGCTTTT CTAGACTTTA AAGAATAAAG TCTTTGGGAG 2460
GTCTCACGGT GTAGAGAGGA GCTTTGAGGC CACCCGCACA AAATTCACCC AGAGGGAAAT 2520
CTCGTCGGAA GGACACTCAC GGCAGTCTG GATCACCTGT GTATGTCAAC AGAAGGGATA 2580
CCGTCTCCTT GAAGAGGAAA CTCTGTCACT CCTCATGCCT GTCTAGCTCA TACACCCATT 2640
TCTCTTTGCT TCACAGGTTT TAACTGGTT TTTTGCATAC TGCTATATAA TTCTCTGTCT 2700
CTCTCTGTTT ATCTCTCCCC TCCCTCCCCT CCCCTTCTTC TCCATCTCCA TTCTTTTGAA 2760
TTTCTCATC CCTCCATCTC AATCCCGTAT CTACGCACCC CCCCCCCCCC AGGCAAAGCA 2820
GTGCTCTGAG TATCACATCA CACAAAAGGA ACAAAGCGA AACACACAAA CCAGCCTCAA 2880
CTTACACTTG GTTACTCAA AGAACAAGAG TCAATGGTAC TTGTCTTAGC GTTTTGGAA 2940
AGGAAAACAG GAACCCACCA AACCAACCAA TCAACCAAC AAAGAAAAAA TTCCACAATG 3000
AAAGAATGTA TTTTGTCTTT TTGCATTTTG GTGTATAAGC CATCAATATT CAGCAAAATG 3060
ATTCCTTTCT TTAATAAAAA AAATGTGGAG GAAAGTAGAA ATTTACCAAG GTTGTGGGCC 3120
CAGGCGCTTA AATTCACAGA TTTTAAAAAC GAGAAAAACA CACAGAAGAA GCTACCTCAG 3180
GTGTTTTTAC CTCAGCACCT TGCTCTGTG TTTCCCTTAG AGATTTTGTA AAGCTGATAG 3240
TTGGAGCATT TTTTATTTT TTTAATAAAA ATGAGTTGGA AAAAAAATAA GATATCAACT 3300
GCCAGCCTGG AGAAGGTGAC AGTCCAAGTG TGCAACAGCT GTTCTGAATT GTCTTCCGCT 3360
AGCCAAGAAC CNATATGGCC TTCTTTTGGA CAAACCTTGA AAATGTTTAT TT 3412

```

<210> 7  
 <211> 1946  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <400> 7

GCTGTAGCGG	AGGGGCTGGG	GGGCTGCTCT	GTCCCCCTTC	TTGCGCGCTG	CGGCCTCAGC	60
CCACCCAGAG	GCCGGGGTGG	GAGGGCGAGT	GCTCAGCTTC	CCGGGTTAGG	AGCCGGAAAA	120
TTCAAATCCG	AAATATTCCA	CCCCAGCTCC	GATGGGAAGT	ACTGGACAGC	CTGCTGGCTC	180
AGTATGGTAC	AGTAGAGAAC	TGTGAGCAAG	TGAACACCGA	GAGTGAGACG	GCAGTGGTGA	240
ATGTACACCTA	TTCCAACCGG	GAGCAGACCA	GGCAAGCCAT	CATGAAGCTG	AATGGCCACC	300
AGTTGGAGAA	CCATGCCCTG	AAGGTCTCCT	ACATCCCCGA	TGAGCAGATA	GCACAGGGAC	360
CTGAGAATGG	GCGCCGAGGG	GGCTTTGGCT	CTCGGGGTCA	GCCCCGCCAG	GGCTCACCTG	420
TGGCAGCGGG	GGCCCCAGCC	AAGCAGCAGC	AAGTGGACAT	CCCCCTTCGG	CTCCTGGTGC	480
CCACCCAGTA	TGTGGGTGCC	ATTATTGGCA	AGGAGGGGGC	CACCATCCGC	AACATCACAA	540
AACAGACCCA	GTCCAAGATA	GACGTGCATA	GGAAAGAGAA	CGCAGGTGCA	GCTGAAAAAG	600
CCATCAGTGT	GCACTCCACC	CCTGAGGGCT	GCTCCTCCGC	TTGTAAGATG	ATCTTGGAGA	660
TTATGCATAA	AGAGGCTAAG	GACACCAAAA	CGGCTGACGA	GGTCCCCCTG	AAGATCCTGG	720
CCCATATAAA	CTTTGTAGGG	CGTCTCATTG	GCAAGGAAGG	ACGGAACCTG	AAGAAGGTAG	780
AGCAAGATAC	CGAGACAAAA	ATCACCATCT	CCTCGTTGCA	AGACCTTACC	CTTTACAACC	840
CTGAGAGGAC	CATCACTGTG	AAGGGGGCCA	TCGAGAATTG	TTGCAGGGCC	GAGCAGGAAA	900
TAATGAAGAA	AGTTCGGGAG	GCCTATGAGA	ATGATGTGGC	TGCCATGAGC	TCTCACCTGA	960
TCCCTGGCCT	GAACCTGGCT	GCTGTAGGTC	TTTTCCCAGC	TCATCCAGC	GCAGTCCCGC	1020
CGCCTCCAG	CAGCGTTACT	GGGGCTGCTC	CCTATAGCTC	CTTTATGCAG	GCTCCCGAGC	1080
AGGAGATGGT	GCAGGTGTTT	ATCCCCGCCC	AGGCAGTGGG	CGCCATCATC	GGCAAGAAGG	1140
GGCAGCACAT	CAAACAGCTC	TCCCGGTTTG	CCAGCGCCTC	CATCAAGATT	GCACCACCCG	1200
AAACACCTGA	CTCCAAAGTT	CGTATGGTTA	TCATCACTGG	ACCGCCAGAG	GCCCAATTCA	1260
AGGCTCAGGG	AAGAATCTAT	GGCAAACCTA	AGGAGGAGAA	CTTCTTTGGT	CCCAAGGAGG	1320
AAGTGAAGCT	GGAGACCCAC	ATACGTGTGC	CAGCATCAGC	AGCTGGCCGG	GTCATTGGCA	1380
AAGGTGGAAA	AACGGTGAAC	GAGTTGCAGA	ATTTGACGGC	AGCTGAGGTG	GTAGTACCAA	1440
GAGACCAGAC	CCCTGATGAG	AACGACCAGG	TCATCGTGAA	AATCATCGGA	CATTTCTATG	1500
CCAGTCAGAT	GGCTCAACGG	AAGATCCGAG	ACATCCTGGC	CCAGGTTAAG	CAGCAGCATC	1560
AGAAGGGACA	GAGTAACCAG	GCCCAGGCAC	GGAGGAAGTG	ACCAGCCCCT	CCCTGTCCCT	1620
TNGAGTCCAG	GACAACAACG	GGCAGAAATC	GAGAGTGTGC	TCTCCCCGGC	AGGCCGTGAG	1680
ATGAGTGGGA	ATCCGGGACA	CNTGGGCCGG	GCTGTAGATC	AGGTTTGCCC	ACTTGATTGA	1740
GAAAGATGTT	CCAGTGAGGA	ACCCTGATCT	NTCAGCCCCA	AACACCCACC	CAATTGGCCC	1800
AACACTGTNT	GCCCCCTCGG	GTGTCAGAAA	TTNTAGCGCA	AGGCACTTTT	AAACGTGGAT	1860
TGTTTTAAAGA	AGCTCTCCAG	GCCCCACCAA	GAGGGTGGAT	CACACCTCAG	TGGGAAGAAA	1920
AATAAAATTT	CCTTCAGGTT	TTAAAA				1946

<210> 8  
 <211> 3283  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <400> 8

GGCAGCGGAG	GAGGCGAGGA	GCGCCGGGTA	CCGGGCCCGG	GGAGCCGCGG	GCTCTCGGGG	60
AAGAGACGGA	TGATGAACAA	GCTTTACATC	GGGAACCTGA	GCCCCGCCGT	CACCGCCGAC	120
GACCTCCGGC	AGCTCTTTGG	GGACAGGAAG	CTGCCCCTGG	CGGGACAGGT	CCTGCTGAAG	180
TCCGGCTACG	CCTTCGTGGA	CTACCCCGAC	CAGAACTGGG	CCATCCGCGC	CATCGAGACC	240
CTCTCGGGTA	AAGTGGAAAT	GCATGGGAAA	ATCATGGAAG	TTGATTACTC	AGTCTCTAAA	300
AAGCTAAGGA	GCAGGAAAAT	TCAGATTCTG	AACATCCCTC	CTCACCTGCA	GTGGGAGGTG	360
TTGGATGGAC	TTTTGGGTCA	ATATGGGACA	GTGGAGAATG	TGGAACAAGT	CAACACAGAC	420
ACAGAAACCG	CCGTTGTCAA	CGTCACATAT	GCAACAAGAG	AAGAAGCAAA	AATAGCCATG	480
GAGAAGCTAA	GCGGGCATCA	GTTTGAGAAC	TACTCCTTCA	AGATTTCCCTA	CATCCCGGAT	540
GAAGAGGTGA	GCTCCCCTTC	GCCCCCTCAG	CGAGCCCAGC	GTGGGGACCA	CTCTTCCCGG	600
GAGCAAGGCC	ACGCCCTTGG	GGGCACTTCT	CAGGCCAGAC	AGATTGATTT	CCCGCTGCGG	660
ATCCTGGTCC	CCACCCAGTT	TGTTGGTGCC	ATCATCGGAA	AGGAGGGCTT	GACCATAAAG	720
AACATCACTA	AGCAGACCCA	GTCCCGGGTA	GATATCCATA	GAAAAGAGAA	CTCTGGAGCT	780
GCAGAGAAGC	CTGTCACCAT	CCATGCCACC	CCAGAGGGGA	CTTCTGAAGC	ATGCCGCATG	840
ATTCTTGAAA	TCATGCAGAA	AGAGGCAGAT	GAGACCAAAC	TAGCCGAAGA	GATTCCTCTG	900
AAAATCTTGG	CACACAATGG	CTTGGTTGGA	AGACTGATTG	GAAAAGAAGG	CAGAAATTTG	960

AAGAAAATTG	AACATGAAAC	AGGGACCAAG	ATAACAATCT	CATCTTTGCA	GGATTTGAGC	1020
ATATACAACC	CGGAAAGAAC	CATCACTGTG	AAGGGCACAG	TTGAGGCCTG	TGCCAGTGCT	1080
GAGATAGAGA	TTATGAAGAA	GCTGCGTGAG	GCCTTTGAAA	ATGATATGCT	GGCTGTTAAC	1140
ACCCACTCCG	GATACTTCTC	CAGCCTGTAC	CCCCATCACC	AGTTTGGCCC	GTTCCCGCAT	1200
CATCACTCTT	ATCCAGAGCA	GGAGATTGTG	AATCTCTTCA	TCCCAACCCA	GGCTGTGGGC	1260
GCCATCATCG	GGAAGAAGGG	GGCACACATC	AAACAGCTGG	CGAGATTCGC	CGGAGCCTCT	1320
ATCAAGATTG	CCCCTGCGGA	AGGCCAGAGC	GTCAGCGAAA	GGATGGTCAT	CATCACCGGG	1380
CCACCGGAAG	CCCAGTTCAA	GGCCCAGGGA	CGGATCTTTG	GGAAACTGAA	AGAGGAAAAC	1440
TTCTTTAACC	CCAAAGAAGA	AGTGAAGCTG	GAAGCGCATA	TCAGAGTGCC	CTCTTCCACA	1500
GCTGGCCGGG	TGATTGGCAA	AGGTGGCAAG	ACCGTGAACG	AACTGCAGAA	CTTAACCACT	1560
GCAGAACTCA	TCGTGCCTCG	TGACCAAACG	CCAGATGAAA	ATGAGGAAGT	GATCGTCAGA	1620
ATTATCGGGC	ACTTCTTTGC	TAGCCAGACT	GCACAGCGCA	AGATCAGGGA	AATTGTACAA	1680
CAGGTGAAGC	AGCAGGAGCA	GAAATACCCCT	CAGGGAGTCG	CCTCACAGCG	CAGCAAGTGA	1740
GGCTCCACAC	GGCACCAGCA	AAACAACGGA	TGAATGTAGC	CCTTCCAACA	CCTGACAGAA	1800
TGAGACCAAA	CGCAGCCAGC	CAGATCGGGA	GCAAACCAAA	GACCATCTGA	GGATGAGAA	1860
GTCTGCGGAG	GCGGCCAGGG	ACTCTGCCGA	GGCCCTGAGA	ACCCCAGGGG	CCGAGGAGGG	1920
GCGGGGAAGG	TCAGCCAGGT	TTGCCAGAAC	CACCGAGCCC	CGCCTCCCGC	CCCCCAGGGC	1980
TTCTGCAGGC	TTTCAAGCATC	CACTTCACCA	TCCACTCGGA	TCTCTCCTGA	ACTCCACGA	2040
CGCTATCCCT	TTTAGTTGAA	CTAACATAGG	TGAACGTGTT	CAAAGCCAAG	CAAAATGCAC	2100
ACCCTTTTTC	TGTGGCAAAT	CGTCTCTGTA	CATGTGTGTA	CATATTAGAA	AGGGAAGATG	2160
TTAAGATATG	TGGCCTGTGG	GTTACACAGG	GTGCCTGCAG	CGGTAATATA	TTTTAGAAAT	2220
AATATATCAA	ATAACTCAAC	TAACTCCAAT	TTTTAATCAA	TTATTAATTT	TTTTTCTTTT	2280
TTAAAGAGAA	AGCAGGCTTT	TCTAGACTTT	AAAGAATAAA	GTCTTTGGGA	GGTCTCACGG	2340
TGTAGAGAGG	AGCTTTGAGG	CCACCCGCAC	AAAATTCACC	CAGAGGGAAA	TCTCGTCGGA	2400
AGGACACTCA	CGGCAGTTCT	GGATCACCTG	TGTATGTCAA	CAGAAGGGAT	ACCGTCTCCT	2460
TGAAGAGGAA	ACTCTGTCAC	TCCTCATGCC	TGTCTAGCTC	ATACACCCAT	TTCTCTTTGC	2520
TTACACAGTT	TTAAACTGGT	TTTTTGCATA	CTGCTATATA	ATTCTCTGTC	TCTCTCTGTT	2580
TATCTCTCCC	CTCCCTCCCC	TCCCCTTCTT	CTCCATCTCC	ATTCTTTTGA	ATTTCTCAT	2640
CCCTCCATCT	CAATCCCGTA	TCTACGCACC	CCCCCCCCCC	CAGGCAAAGC	AGTGCTCTGA	2700
GTATCACATC	ACACAAAAGG	AACAAAAGCG	AAACACACAA	ACCAGCCTCA	ACTTACACTT	2760
GGTTACTCAA	AAGAACAAGA	GTCAATGGTA	CTTGTCTTAG	CGTTTTGGAA	GAGGAAAACA	2820
GGAACCCACC	AAACCAACCA	ATCAACCAAA	CAAAGAAAAA	ATTCCACAAT	GAAAGAATGT	2880
ATTTTGTCTT	TTTGCATTTT	GGTGTATAAG	CCATCAATAT	TCAGCAAAAT	GATTCCTTTC	2940
TTTAAAAAAA	AAAATGTGGA	GGAAAGTAGA	AATTTACCAA	GGTTGTTGGC	CCAGGGCGTT	3000
AAATTCACAG	ATTTTTTTTAA	CGAGAAAAAC	ACACAGAAGA	AGCTACCTCA	GGTGTTTTTTA	3060
CCTCAGCACC	TTGCTCTTGT	GTTTCCCTTA	GAGATTTTGT	AAAGCTGATA	GTTGGAGCAT	3120
TTTTTTTATTT	TTTTAATAAA	AATGAGTTGG	AAAAAAAATA	AGATATCAAC	TGCCAGCCTG	3180
GAGAAGGTGA	CAGTCCAAGT	GTGCAACAGC	TGTTCTGAAT	TGTCTTCCGC	TAGCCAAGAA	3240
CCNATATGGC	CTTCTTTTGG	ACAAACCTTG	AAAATGTTTA	TTT		3283

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/05766

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : G01N 33/574, 33/48; C12Q 1/68

US CL : 435/7.23, 6; 436/64

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/7.23, 6; 436/64

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	TURECI, O. et al. Expression of SSX Genes in Human Tumors International Journal of Cancer. 1998, Vol. 77, No. 1, pages 19-23, especially page 19.	41-52
A	GURE, A.O. et al. SSX: A Multigene Family with Several Members Transcribed in Normal Testis and Human Cancer. International Journal of Cancer. 1997, Vol. 72, No. 6, pages 965-971, especially page 965.	41-52
A	CHEN, et al. A Testicular Antigen Aberrantly Ex[pressed in Human Cancers Detected by Autologous Antibody Screening. Proceedings of the National Academy of Science. March 1997, Vol. 94, No. 5, pages 1914-1918, especially page 1914.	41-52



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

21 JULY 1999

Date of mailing of the international search report

11 AUG 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

YVONNE EYLER

Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/05766

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	CHEN, et al. Identification of Multiple Cancer/Testis Antigens by Allogeneic Antibody Screening of a Melanoma Cell Line Library" Proceedings of the National Academy of Science. June 1998, Vol. 95, No. 12, pages 6919-6923, especially page 6919.	41-52
P,Y	UGUR, S. et al. Expression of Multiple Cancer/Testis (CT) Antigens in Breast Cancer and Melanoma: Basis for Polyvalent CT Vaccine Strategies. International Journal of Cancer. 1998, Vol. 78, No. 3, pages 387-389, especially page 387.	41-52

**THIS PAGE IS BLANK**